



Diet of three insectivorous birds in Nagapattinam District, Tamil Nadu, India – a preliminary study

S. Asokan¹, A. Mohamed Samsoor Ali² & R. Manikannan³

¹Reader & ³Ph.D. Research Scholar, Department of Zoology & Division of Wildlife Biology, A.V.C. College (Autonomous), Mannampandal, Mayiladuthurai, Tamil Nadu 609305, India

²Senior Research Fellow, Owl Research Project, Department of Zoology, Saraswathi Narayanan College (Autonomous), Perungudi, Madurai, Tamil Nadu 625022, India

Email: beesasokan@yahoo.co.in¹; amsamsoor@yahoo.co.in² (corresponding author); manikannan@yahoo.co.in³

Date of online publication 26 June 2009
ISSN 0974-7907 (online) | 0974-7893 (print)

Editor: C. Srinivasulu

Manuscript details:

Ms # o2145
Received 13 February 2009
Final received 08 May 2009
Finally accepted 30 May 2009

Citation: Asokan, S., A.M.S. Ali & R. Manikannan (2009). Diet of three insectivorous birds in Nagapattinam District, Tamil Nadu, India – a preliminary study. *Journal of Threatened Taxa* 1(6): 327-330.

Copyright: © S. Asokan, A. Mohamed Samsoor Ali & R. Manikannan 2009. Creative Commons Attribution 3.0 Unported License. JoTT allows unrestricted use of this article in any medium for non-profit purposes, reproduction and distribution by providing adequate credit to the authors and the source of publication.

Acknowledgements: We are highly grateful to the Ministry of Environment and Forests, Govt. of India for their financial support (14/14/2001/ERS/RE) to carry out the above work in Nagapattinam District, Tamil Nadu. We wish to express our sincere thanks to the HOD and other staff members of zoology and the Principal and Management of A.V.C. College (Autonomous), Mannampandal for having rendered facilities and encouragement. We are thankful to M/s. P. Radhakrishnan and M. Chakravarthi for their field help.

Abstract: The dietary composition of the White-breasted Kingfisher *Halcyon smyrnensis*, the Small Bee-eater *Merops orientalis* and the Black Drongo *Dicrurus macrocercus* was studied between 2005 and 2006 in Nagapattinam District, Tamil Nadu, India by analyzing regurgitated pellets. The analysis revealed that the White-breasted Kingfisher preys mainly on arthropods (83.40%) and less on vertebrates; seven orders of insects were identified, with Coleoptera, Hemiptera, Hymenoptera and Orthoptera predominant. The small bee-eater diet is composed of Coleoptera (22.3%), Hymenoptera (20.8%), Hemiptera (14.1%), Orthoptera (12.6%), Odonata (10.7%), Lepidoptera (10.4%) and Diptera (8.6%). Beetles were also found to be the most frequent prey (23.7%) in the diet of black drongos, followed by Hemiptera (21.6%), Orthoptera (19.3%), Hymenoptera (14.4%), Lepidoptera (7.5%), Diptera (6.8%) and Odonata (6.0%).

Keywords: Bio-control, birds, Coleoptera, diet, Hemiptera, Hymenoptera, Insectivory

INTRODUCTION

Agricultural ecosystem provides a concentrated and highly predictable source of food to many birds in the form of grain, seeds, insects and rodents. A large number of birds directly or indirectly depend upon crop fields, making them integral parts of agro-ecosystems. Insectivorous birds are beneficial to agriculturists since they keep a check on populations of various insect pests of crops. As enemies of insects, birds stand supreme among vertebrates because they are highly mobile and are able to congregate quickly in large numbers when sudden outbreaks of insect pests occur (Dhindsa & Saini 1994). Thus birds are highly motivated, efficient and cost-effective insect pest controllers.

Detailed analysis of food and feeding behaviour of insectivorous birds is fundamental to assessments of their economic importance. However, information on the feeding ecology of Indian birds is limited, being largely based on the findings of Beresford (1944), Mukherjee (1975), Mathew et al. (1978) and Ali & Ripley (1983). In southern India, reports on the food and feeding ecology of insectivorous birds are few and scanty (Mathew et al. 1978; Natarajan 1993; Asokan 1998; Sivakumaran & Thiyagesan 2003; Asokan et al. 2008). The present paper deals with the diets of three common insectivorous birds: the White-breasted Kingfisher *Halcyon smyrnensis*, Small Bee-eater *Merops orientalis* and Black Drongo *Dicrurus macrocercus* in Nagapattinam District, Tamil Nadu, India. Beyond general observations on their distribution and breeding seasons, our knowledge on feeding ecology of these three species is limited.

MATERIALS AND METHODS

The present study was conducted in two village's viz., Thiruvananthapuram and Manakudi (18°18'N & 79°50'E) of Cauvery Delta in Nagapattinam District, Tamil Nadu, India between 2005 and 2006. The study area is a portion of the granary of southern India because of extensive cultivation of paddy, sugarcane, cotton, groundnut, banana, pulses and other cereals. The river Cauvery and its tributaries, namely, *Kollidam*, *Uppanar*, *Vellar*, *Manjalar* and *Arasalar* are the major perennial water sources in the study area. The north-east monsoon brings rain between October and December (amounting to 65% of the total rainfall in a year).

Analysis of regurgitated pellets is a reliable, non-invasive method for studying food habits of insectivorous birds (Ralph et al. 1985). The regurgitated pellets were collected opportunistically at perch and roost-sites around the agricultural lands. The collected pellets were bagged, labeled and dried in a hot air oven at 60°C for identification of prey remains (Herrera & Ramirez 1974; Ralph et al. 1985). The prey fragments were identified



OPEN ACCESS | FREE DOWNLOAD

Table 1. Frequency and percent composition of various prey items in the diet of the White-breasted Kingfisher (n=638 pellets)

Prey items	Frequency	% composition	Rank
Invertebrates			
Arthropoda			
Coleoptera (beetles)	1290	21.95	I
Hemiptera (bugs)	1009	17.17	II
Hymenoptera (ants, bees and wasps)	866	14.74	III
Orthoptera (grasshoppers and crickets)	806	13.72	IV
Diptera (flies)	411	6.99	V
Odonata (dragonflies)	279	4.75	VII
Lepidoptera (butterflies and moths)	240	4.08	VIII
Total	4901	83.4	
Mollusca			
Crab	372	6.33	VI
Vertebrates			
Pisces	166	2.83	X
Amphibians	195	3.32	IX
Reptiles	165	2.81	XI
Unidentified prey items	77	1.31	XII
Total	5876	100	

Table 3. Frequency and percent composition of various prey items in the diet of the Small Bee-eater (n=714 pellets)

Prey items	Frequency	% composition	Rank
Coleoptera (beetles)	1950	22.3	I
Hemiptera (bugs)	1230	14.1	III
Hymenoptera (ants, bees and wasps)	1821	20.8	II
Orthoptera (grasshoppers and crickets)	1102	12.6	IV
Diptera (flies)	757	8.6	VII
Odonata (dragonflies)	933	10.7	V
Lepidoptera (butterflies and moths)	914	10.4	VI
Unidentified insects	47	0.5	VIII
Total	8754	100	

using a dissecting microscope and reference books. We identified the prey taxa by means of analyzing various body-parts in the remains of the diets as follows: Coleoptera (mandibles, elytra and leg fragments), Hemiptera (H-shaped tergal plates, mouthparts and leg fragments), Hymenoptera (mouthparts, leg fragments and wing fragments), Orthoptera

(raptorial leg fragments and mandibles), Diptera (antennae, eyes and wings), Odonata (wings, leg fragments and head capsules), Lepidoptera (wing scales and proboscis), crabs (pedipalps and shell fragments), fish (cartilage bones and scales), amphibians (mandibles and pelvic girdle) and reptiles (mandibles).

RESULTS AND DISCUSSION

Analysis of the diet of the White-breasted Kingfisher based on 638 regurgitated pellets reveals that it preys primarily on arthropods (83.40 %). Among the insects Coleoptera (21.95 %), Hemiptera (17.17 %), Hymenoptera (14.74 %) and Orthoptera (13.72 %) occur most frequently in its diet (Table 1). In addition, crabs, fishes, amphibians and reptiles comprise about 16.60% of the total diet (Table 1). Our findings are consistent with the results of some of the earlier studies (e.g., Sen 1944; Mukherjee 1975; Yahya 2001). In another study carried out in Malaysia (Burton 1998), insects - mostly grasshoppers, ants and beetles - and lizards were observed to be the main diet of White-breasted Kingfisher. A similar trend was found in terms of prey composition, with as many as 613 of the total 638 pellets (96%) having remains of insects belonging to order Coleoptera, followed by Hemiptera (93%), Orthoptera (91%) and Hymenoptera (89%). Other prey remains were found in less than 50% of the total pellets analyzed (Table 2). The remains of fishes, mostly restricted to rainy season, were detected in 103 pellets. During the rainy season (September to January) when the water level is much higher and the river and ponds largely inundated, the birds fed more on aquatic animals like fish and frogs, while during the drier months from February to August they consumed more terrestrial organisms.

The remains of 8757 insect prey items were found in the regurgitated pellets of Small Bee-eaters. The most dominant prey were Coleoptera (22.3%), Hymenoptera (20.8%), Hemiptera (14.1%), Orthoptera (12.6%), Odonata (10.7%), Lepidoptera (10.4%) and Diptera (8.6%) (Table 3). Earlier, Asokan (1998) found that beetles and Hymenoptera were the principal food items of Small Bee-eaters in Nagapattinam District. Mathew et al. (1978) reported that they consumed almost equal proportions of Hymenoptera, Coleoptera, Odonata and Lepidoptera. However, Fry (1984) who studied the Little Bee-eater *Merops pusillus* in Africa, reported that Hymenoptera formed the major diet. The percentage occurrence of Hymenoptera in the study area in the diet of Small Bee-eaters

Table 2. Comparative statistics for different prey items found in the White-breasted Kingfisher pellets (n=638)

	Coleoptera	Hemiptera	Hymenoptera	Orthoptera	Diptera	Odonata	Lepidoptera	Crab	Pisces	Amphibians	Reptiles
Presence	613	596	569	584	302	188	161	298	103	182	153
Presence %	96%	93%	89%	91%	47%	29%	25%	47%	16%	28%	24%
Absence	25	42	69	54	336	450	477	340	535	456	485
Absence %	4%	7%	11%	9%	53%	71%	75%	53%	84%	72%	76%
Minimum	0	0	0	0	0	0	0	0	0	0	0
Maximum	19	8	6	11	2	4	2	4	4	2	2
Range	0-19	0-8	0-6	0-11	0-2	0-4	0-2	0-4	0-4	0-2	0-2
Sum	1290	1009	866	806	411	279	240	372	166	195	165

Presence - The number of pellets containing remains of the respective prey item; Presence % - The percentage of pellets containing remains of the respective prey item; Absence - The number of pellets not containing remains of the respective prey item; Absence % - The percentage of pellets not containing remains of the respective prey item; Minimum - The minimum number of remains in any pellets of the respective prey item; Maximum - The maximum number of remains in any pellets of the respective prey item; Range - The range of remains in pellets of the respective prey item; Sum - The total number of remains of the respective prey item in all pellets.

Table 4. Comparative statistics for different prey items found in the Small Bee-eater pellets (n=714)

	Coleoptera	Hemiptera	Hymenoptera	Orthoptera	Diptera	Odonata	Lepidoptera
Presence	688	496	661	579	368	515	420
Presence %	96%	69%	93%	81%	52%	72%	59%
Absence	26	218	53	135	346	199	294
Absence %	4%	31%	7%	19%	48%	28%	41%
Minimum	0	0	0	0	0	0	0
Maximum	18	5	6	10	2	4	3
Range	0-18	0-5	0-6	0-10	0-2	0-4	0-3
Sum	1950	1230	1821	1102	757	933	914

Presence - The number of pellets containing remains of the respective prey item; Presence % - The percentage of pellets containing remains of the respective prey item; Absence - The number of pellets not containing remains of the respective prey item; Absence % - The percentage of pellets not containing remains of the respective prey item; Minimum - The minimum number of remains in any pellets of the respective prey item; Maximum - The maximum number of remains in any pellets of the respective prey item; Range - The range of remains in pellets of the respective prey item; Sum - The total number of remains of the respective prey item in all pellets.

Table 5. Frequency and percent composition of various prey items in the diet of the Black Drongo (n=676 pellets)

Prey items	Frequency	% composition	Rank
Coleoptera (beetles)	2166	23.7	I
Hemiptera (bugs)	1971	21.6	II
Hymenoptera (ants, bees and wasps)	1315	14.4	IV
Orthoptera (grasshoppers and crickets)	1766	19.3	III
Diptera (flies)	619	6.8	VI
Odonata (dragonflies)	548	6	VII
Lepidoptera (butterflies and moths)	682	7.5	V
Unidentified insects	61	0.7	VIII
Total	9128	100	

could be attributed to less numbers of bee-hives in the study area. In terms of prey composition, most pellets had remains of insects belonging to order Coleoptera (96%) followed by Hymenoptera (93%), Orthoptera (81%) and Odonata (72%). The other prey remains such as Hemiptera, Diptera and Lepidoptera were found in less than 60% of pellets (Table 4).

The analysis of 676 pellets of Black Drongo indicated the high frequency occurrence of insects belonging to order Coleoptera (23.7%) followed by Hemiptera (21.6%), Orthoptera (19.3%), Hymenoptera (14.4%), Lepidoptera (7.5%), Diptera (6.8%) and Odonata (6.0%) (Table 5). Mathew et al. (1978) reported that Coleoptera, Hemiptera and Orthoptera as major food of Black Drongos, and the present study showed a similar trend. Hussain & Bhalla (1937) found that Black Drongos feed

largely on grasshoppers and locusts. In terms of prey composition totally 676 regurgitated pellets were analyzed, of these 92% of the pellets contained Coleopteran remains, 83% contained Hemipteran remains, 76% contained Orthopteran remains and 72% contained Hymenopteran remains. The Diptera, Odonata and lepidoptera were found in 368 (54%), 389 (58%) and 413 (61%) pellets respectively (Table 6).

White-breasted Kingfishers, Small Bee-eaters and Black Drongos consumed almost similar proportions of Coleopterans, Hemipterans, Hymenopterans and Orthopterans in our study. These three species were predominantly found in agricultural lands and the above insect prey formed principal food items due to their greater availability. Moorman et al. (2007) reported that Coleopterans and Hemipterans formed the major diet of foliage-gleaning and ground-gleaning insectivorous birds. Many other investigators also have determined Coleopterans, Hemipterans and Hymenopterans (Mason & Maxwell-Lefroy 1912; Sodhi 1986; Kaspari & Joern 1993; Parasharya et al. 1994; Sivakumaran & Thiyagesan 2003; Yard et al. 2004; Asokan et al. 2008) as primary food resource of insectivorous birds.

The findings of the present study revealed that a wide variety of insects are consumed by the three birds showed and hence they act as very active bio-control agents against agricultural insects pests. Nathan & Rajendran (1982) reported that Black Drongos destroying injurious insects like stem-borer, skippers and leaf rollers in enormous numbers in the rice ecosystem of Pondicherry region. Parasharya et al. (1994)

Table 6. Comparative statistics for different prey items found in the Black Drongo pellets (n=676)

	Coleoptera	Hemiptera	Hymenoptera	Orthoptera	Diptera	Odonata	Lepidoptera
Presence	624	562	485	517	368	389	413
Presence %	92%	83%	72%	76%	54%	58%	61%
Absence	52	114	191	159	308	287	263
Absence %	8%	17%	28%	24%	46%	42%	39%
Minimum	0	0	0	0	0	0	0
Maximum	18	5	5	11	3	4	3
Range	0-18	0-5	0-5	0-11	0-3	0-4	0-3
Sum	2166	1971	1315	1766	619	548	682

Presence - The number of pellets containing remains of the respective prey item; Presence % - The percentage of pellets containing remains of the respective prey item; Absence - The number of pellets not containing remains of the respective prey item; Absence % - The percentage of pellets not containing remains of the respective prey item; Minimum - The minimum number of remains in any pellets of the respective prey item; Maximum - The maximum number of remains in any pellets of the respective prey item; Range - The range of remains in pellets of the respective prey item; Sum - The total number of remains of the respective prey item in all pellets.

also stated that White-breasted Kingfishers, Small Bee-eaters and Black Drongos are important bio-control agents against white grub (*Holotricha* sp.), an important subterranean pest damaging root system of several crops.

Interestingly, the local farmers are aware of the beneficial role of these insectivorous birds in bio-control of crop pests. Farmers can encourage these birds in agricultural fields by providing perching sites, by keeping dried tree stumps in different localities in the rice fields, and also by planting plantain along the farm boundaries.

REFERENCES

- Ali, S. & S.D. Ripley (1983). *The Book of Indian Birds*. Oxford University Press, London.
- Asokan, S. (1998). Food and feeding habits of the Small Green Bee-eater *Merops orientalis* in Mayiladuthurai. *Journal of Ecobiology* 10(3): 199-204.
- Asokan, S., A.M.S Ali, R. Manikannan & P. Radhakrishnan (2008). Studies on insect composition in the diet of Common Myna *Acridotheres tristis* L. *Trends in Kalis Research* 3(1): 72-79.
- Beresford, G. (1944). The winter food of birds in Kashmir. *Journal of the Bombay Natural History Society* 45: 86-88.
- Burton, N.H.K. (1998). Notes on the diet of nestling White-throated Kingfishers *Halcyon smyrnensis* in Malaysia. *Forktail* 14: 79-80.
- Dhindsa, M.S. & H.K. Saini (1994). Agricultural ornithology: an Indian perspective. *Journal of Bioscience* 19: 391-402.
- Fry, C.H. (1984). *The bee-eaters*. Tand A.D. Poyser, Carlton, UK.
- Herrera, C.M. & A. Ramirez (1974). Food of Bee-eaters in Southern Spain. *British Birds* 67: 159-164.
- Hussain, M.A. & H.R. Bhalla (1937). The bird enemies of the cotton roller *Svlepta degogata* at Khanewal Multan (Punjab). *Indian Journal of Agricultural Science* 7: 785-792.
- Kaspari, M. & A. Joern (1993). Prey choice by three insectivorous grassland birds: reevaluating opportunism. *Oikos* 68: 414-430.
- Mason, C.W. & H. Maxwell-Lefroy (1912). The food of birds in India. *Memoris Department of Agriculture in India, Entomological series*, Vol. 3.
- Mathew, D.N., T.C. Narendran & V.J. Zacharias (1978). A comparative study of the feeding habits of certain species of Indian birds affecting agriculture. *Journal of the Bombay Natural History Society* 75(2): 1178-1197.
- Moorman, C.E., L.T. Bowen, J.C. Kilgo, C.E. Sorenson, J.L. Hanula, S. Horn & M.D. Ulyshen (2007). Seasonal diets of insectivorous birds using canopy gaps in a bottomland forest. *Journal of Field Ornithology* 78: 11-20.
- Mukherjee, A.K. (1975). Food-habits of water-birds of the Sundarban, 24 Paraganas District, West Bengal. India – V. *Journal of the Bombay Natural History Society* 72: 85-109.
- Natarajan, V. (1993). Food and feeding habits of Southern Crow-pheasant *Centropus sinensis parroti* Stresemann (Aves: Cuculidae) at Point Calimere, Tamil Nadu. *Journal of the Bombay Natural History Society* 90: 11-16.
- Nathan, S.P.C. & B. Rajendran (1982). Bird fauna of the rice crop ecosystem in Pondicherry region. *Journal of the Bombay Natural History Society* 79: 204-205.
- Parasharya, B.M., J.F. Dodia, K.L. Mathew & D.N. Yadav (1994). Natural regulation of white grub (*Holotricha* sp. Scarabidae) by birds in agro-ecosystem. *Journal of Bioscience* 19: 381-390.
- Ralph, C.P., S.E. Nagata & C.J. Ralph (1985). Analysis of droppings to describe diets of small birds. *Journal of Field Ornithology* 56 (2): 165-174.
- Sen, S.N. (1944). Food of the White-breasted Kingfisher *Halcyon smyrnensis*. *Journal of Bombay Natural History Society* 44:475.
- Sivakumaran, N. & K. Thiyagesan (2003). Population, diurnal activity patterns and feeding ecology of the Indian Roller *Coracias benghalensis*. *Zoos'Print Journal* 18(5): 1091-1095.
- Sodhi, N.S. (1986). Feeding ecology of Indian Pond Heron and its comparison with that of Little Egret. *Pavo* 24: 97-116.
- Yahya, H.S. (2001). Feeding habits of the White-breasted Kingfisher *Halcyon smyrnensis* (Linn.) Family Alcedinidae. *Journal of the Bombay Natural History Society* 98:450-451.
- Yard, H.K., C. Van Riper III, B.T. Brown & M.J. Kearsley (2004). Diets of insectivorous birds along the Colorado River in Grand Canyon, Arizona. *Condor* 106: 106-115.

Author Details: Dr. S. ASOKAN is currently teaching students, undertaking and supervising various research activities. He has produced 7 PhD and 40 MPhil students in various field of Wildlife Biology and Zoology. He has published more than 30 research papers in leading national and international journals. A. MOHAMED SAMSOOR is currently working as a Senior Research Fellow (SRF) in the Owl Research Project (funded by MoEF, Govt. of India) at Saraswathi Narayanan College (Autonomous), Madurai. He graduated with a Master of Science in Wildlife Biology from the A.V.C. College (Autonomous), Mayiladuthurai in 2003 and has been working on the various filed of ornithology since 2004. R. MANIKANNAN is working on diversity of wetland birds in Point Calimere Wildlife Sanctuary for a PhD degree.

Author Contribution: The field study was conducted AMSA and RM under the supervision of Dr. S.A. AMSA analyzed data and wrote the manuscript and Dr. S.A. provided necessary suggestions during manuscript preparation.

