



The Journal of Threatened Taxa (JoTT) is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at www.threatenedtaxa.org. All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use, reproduction, and distribution of articles in any medium by providing adequate credit to the author(s) and the source of publication.

Journal of Threatened Taxa

Building evidence for conservation globally

www.threatenedtaxa.org

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

NOTE

CIRSIUM WALLICHII DC. (ASTERACEAE): A KEY NECTAR SOURCE OF BUTTERFLIES

Bitupan Boruah, Amit Kumar & Abhijit Das

26 October 2020 | Vol. 12 | No. 14 | Pages: 17049–17056

DOI: [10.11609/jott.6008.12.14.17049-17056](https://doi.org/10.11609/jott.6008.12.14.17049-17056)



For Focus, Scope, Aims, Policies, and Guidelines visit <https://threatenedtaxa.org/index.php/JoTT/about/editorialPolicies#custom-0>

For Article Submission Guidelines, visit <https://threatenedtaxa.org/index.php/JoTT/about/submissions#onlineSubmissions>

For Policies against Scientific Misconduct, visit <https://threatenedtaxa.org/index.php/JoTT/about/editorialPolicies#custom-2>

For reprints, contact [<ravi@threatenedtaxa.org>](mailto:ravi@threatenedtaxa.org)

The opinions expressed by the authors do not reflect the views of the Journal of Threatened Taxa, Wildlife Information Liaison Development Society, Zoo Outreach Organization, or any of the partners. The journal, the publisher, the host, and the partners are not responsible for the accuracy of the political boundaries shown in the maps by the authors.

Member



Publisher & Host





Cirsium wallichii DC. (Asteraceae): a key nectar source of butterflies

Bitupan Boruah¹ , Amit Kumar² & Abhijit Das³ ^{1,2,3} Wildlife Institute of India, Chandrabani, Dehradun, Uttarakhand 248002, India.¹ bitupan.kaz@gmail.com (corresponding author), ² amit@wii.gov.in, ³ abhijit@wii.gov.in

In general, both larvae and adult butterflies depend on plant resources (Kitahara et al. 2008; Nimbalkar et al. 2011). Adult butterflies forage on a wide variety of plant species for floral nectar (Courtney 1986; Raju et al. 2004). Butterflies, however, do not collect nectar extensively from all the available flowers (Kunte 2000). Thus, the diversity of the butterfly community of a region is associated with the availability of host plants (Murphy & Wilcox 1986; Kitahara et al. 2008). Also, the diversity and abundance of pollinators such as butterflies are crucial for the reproductive success of flowering plants (Mukherjee et al. 2015). Several wild plants considered as weeds serve as important nectar sources for butterflies (Mukherjee et al. 2015; Kapkoti et al. 2016). One such wild weed, *Cirsium* Mill. (Thistle) of the family Asteraceae has been well recognized as a nectar source of butterflies (Robertson 1928; Tooker et al. 2002; Kapkoti et al. 2016). *Cirsium* is a speciose genus of Asteraceae, with about 200 species distributed in Europe, Asia, North & Central America, and northern Africa (Mabberley 2008; Sahli et al. 2017). Among the species of this genus known from India, *Cirsium wallichii* DC. has been extensively used as a traditional medicinal plant in the Himalaya (Uniyal et al. 2011). Interestingly,

owing to a lack of information on *Cirsium wallichii* DC. as a nectar source of butterflies, the current communication aims to address the value of Wallichii's Thistle not only as a weed, but also as a nectar source of butterflies.

The present study was conducted from May to August, 2019 in Benog Wildlife Sanctuary (30.467°N & 78.027°E), Mussoorie, Uttarakhand, India. The sanctuary is characterized by Banj Oak *Quercus leucotrichophora* forests, Chirpine *Pinus roxburghii* forests and grasslands (Champion & Seth 1968) which harbour at least 335 species of vascular plants (Kumar et al. 2012). The survey was done between 08.00h and 11.00h to record the butterfly species visiting *Cirsium wallichii*. We photographed representatives of each butterfly species from the area. Based on the photographs, identification of the species was carried out using Evans (1932) and Kehimkar (2016).

Cirsium wallichii grows along open and modified stream habitats in the sanctuary as well as near human settlements and agricultural lands at the peripheral area (Image 1A). Leaves are stalkless and pinnately lobed with long spines at the margin. The plant blooms from May–July. Capitula are many-flowered, solitary or clustered and borne on leafless stalks. They are 2–3.4cm

Editor: A.J. Solomon Raju, Andhra University, Visakhapatnam, India.**Date of publication:** 26 October 2020 (online & print)**Citation:** Boruah, B., A. Kumar & A. Das (2020). *Cirsium wallichii* DC. (Asteraceae): a key nectar source of butterflies. *Journal of Threatened Taxa* 12(14): 17049–17056. <https://doi.org/10.11609/jott.6008.12.14.17049-17056>**Copyright:** © Boruah et al. 2020. Creative Commons Attribution 4.0 International License. JoTT allows unrestricted use, reproduction, and distribution of this article in any medium by providing adequate credit to the author(s) and the source of publication.**Funding:** None.**Competing interests:** The authors declare no competing interests.भारतीय वन्यजीव संस्थान
Wildlife Institute of India**Acknowledgements:** The authors wish to thank Uttarakhand Forest Department for necessary field support and permission to conduct the study. We are also thankful to Mussoorie Forest Division for their support specially Dr. Shipra Sharma, Range Forest Officer and Forest staff of Benog Wildlife Sanctuary for helping WII team during the field work. We would like to acknowledge Director and Dean, Wildlife Institute of India, Dehradun for institutional support. Deb S. Goswami and Swati Nawani are also acknowledged for their help during the field work.

Table 1. List of butterfly species foraging on *Cirsium wallichii*

	Scientific name	Common name
A.	Family: Papilionidae	
1.	<i>Graphium sarpedon</i> (Linnaeus, 1758)	Common Bluebottle
2.	<i>Graphium cloanthus</i> (Westwood, 1841)	Glassy Bluebottle
3.	<i>Graphium agamemnon</i> (Linnaeus, 1758)	Tailed Jay
4.	<i>Papilio protenor</i> Cramer, [1775]	Spangle
5.	<i>Papilio bianor</i> Cramer, [1777]	Common Peacock
6.	<i>Papilio polytes</i> Linnaeus, 1758	Common Mormon
B.	Family: Pieridae	
7.	<i>Aporia agathon caphusa</i> (Moore, 1872)	Garhwal Great Blackvein
8.	<i>Aporia agathon agathon</i> (Gray, 1831)	Nepalese Great Blackvein
9.	<i>Aporia leucodice</i> (Eversmann, 1843)	Himalayan Blackvein
10.	<i>Colias erate</i> (Esper, 1805)	Pale Clouded Yellow
11.	<i>Colias fieldii</i> Ménétriér, 1855	Dark Clouded Yellow
12.	<i>Pieris brassicae</i> (Linnaeus, 1758)	Large Cabbage White
13.	<i>Pieris canidia</i> (Linnaeus, 1768)	Indian Cabbage White
14.	<i>Gonepteryx rhamni</i> Linnaeus, 1758	Common Brimstone
15.	<i>Pontia daplidice</i> (Linnaeus, 1758)	Bath White
16.	<i>Belenois aurota</i> (Fabricius, 1793)	Pioneer
C.	Family: Lycaenidae	
17.	<i>Helipophorus sena</i> (Kollar, [1844])	Sorrel Sapphire
18.	<i>Spindasis nipalicus</i> (Moore, 1884)	Silver-grey Silverline
19.	<i>Rapala selira</i> (Moore, 1874)	Himalayan Red Flash
20.	<i>Rapala varuna</i> (Horsfield, [1829])	Indigo Flash
21.	<i>Rapala manea</i> (Hewitson, 1863)	Slate Flash
22.	<i>Aricia agestis</i> (Denis & Schiffermüller, 1775)	Orange-bordered Argus
23.	<i>Lycaena phlaeas</i> (Linnaeus, 1761)	Common Copper
24.	<i>Lampides boeticus</i> (Linnaeus, 1767)	Pea Blue
25.	<i>Chilades pandava</i> (Horsfield, [1829])	Plains Cupid
26.	<i>Celastrina huegelii</i> (Moore, 1882)	Large Hedge Blue
27.	<i>Deudorix epijarbas</i> (Moore, [1858])	Cornelian
D.	Family: Nymphalidae	
28.	<i>Vanessa indica</i> Herbst, 1794	Red Admiral
29.	<i>Vanessa cardui</i> Linnaeus, 1758	Painted Lady
30.	<i>Kaniska canace</i> Linnaeus, 1763	Blue Admiral

	Scientific name	Common name
31.	<i>Aglaia caschmirensis</i> Kollar, 1844	Indian Tortoiseshell
32.	<i>Callerebia annada caeca</i> Moore, 1857	Ringed Argus
33.	<i>Callerebia hybrida</i> Butler, 1880	Hybrid Argus
34.	<i>Callerebia nirmala</i> Moore, 1865	Common Argus
35.	<i>Argynnis hyperbius</i> (Linnaeus, 1763)	Indian Fritillary
36.	<i>Ypthima nareda</i> Kollar, 1844	Large Three-Ring
37.	<i>Ypthima nikaea</i> Moore, 1874	Moore's Five-Ring
38.	<i>Parantica aglea</i> (Stoll, [1782])	Glassy Tiger
39.	<i>Tirumala limniace</i> (Cramer, [1775])	Blue Tiger
40.	<i>Tirumala septentrionis</i> (Butler, 1874)	Dark Blue Tiger
41.	<i>Danaus genutia</i> (Cramer, [1779])	Striped Tiger
42.	<i>Danaus chrysippus</i> (Linnaeus, 1758)	Plain Tiger
43.	<i>Euploea mulciber</i> (Cramer, [1777])	Striped Blue Crow
44.	<i>Argynnis childreni</i> Gray, 1831	Large Silver stripe
45.	<i>Libythea lepita</i> Moore, [1858]	Common Beak
46.	<i>Lasiommata schakra</i> Kollar, 1844	Common Wall
47.	<i>Acraea issoria</i> (Hübner, [1819])	Yellow Coster
48.	<i>Cyrestis thyodamas</i> Doyère, [1840]	Common Map
49.	<i>Junonia iphita</i> Cramer, 1779	Chocolate Pansy
E.	Family: Hesperidae	
50.	<i>Seseria dohertyi</i> Watson, 1893	Himalayan White Flat
51.	<i>Potanthus dara</i> (Kollar, [1844])	Himalayan Dart
52.	<i>Celaenorrhinus leucocera</i> (Kollar, [1844])	Common Spotted Flat
53.	<i>Lobocla liliana</i> Atkinson, 1871	Marbled Flat
54.	<i>Celaenorrhinus dhanada</i> (Moore, [1866])	Himalayan Yellow-banded Flat
55.	<i>Pseudocoladenia dan</i> (Fabricius, 1787)	Fulvous Pied Flat
56.	<i>Tagiades menaka</i> Moore, 1865	Spotted Snow Flat
57.	<i>Celaenorrhinus munda</i> Moore, 1884	Himalayan Spotted Flat
58.	<i>Aeromachus stigmata</i> Moore, 1878	Veined Scrub Hopper
59.	<i>Notocrypta feisthamelii</i> Boisduval, 1832	Spotted Demon
60.	<i>Pedesta masuriensis</i> Moore, 1878	Mussoorie Bush Bob
61.	<i>Polytremis discreta</i> (Elwes & Edwards, 1897)	Himalayan Swift
62.	<i>Parnara</i> sp.	Swift sp.



Image 1. *Cirsium wallichii*: A—habit | B—inflorescence | C—flower. © Bitupan Boruah.

across, homogamous, bisexual, discoid, and clustered in corymbose racemes (Image 1B). Florets are about 2cm long, pale-white, corolla tube long, limb five-toothed and pappus hair pale-white. Outer involucre bracts are lanceolate with spreading erect or recurved spines; inner bracts dilated, lanceolate-ovate and incurved near the apex (Image 1C).

During recent field explorations in the Benog Wildlife Sanctuary, a total of 62 species and subspecies of butterflies belonging to 45 genera and five families foraging on *Cirsium wallichii* for nectar were documented (Table 1 and Images 2–5). The species assemblage includes Nymphalidae (35.5%), Hesperidae (22.6%), Lycaenidae (17.7%), Pieridae (16.1%) and Papilionidae (9.7%). Among the recorded butterflies, five species such as *Aporia agathon*, *Gonepteryx rhamni*, *Celaenorrhinus munda*, *Vanessa cardui*, and *Vanessa indica* frequently visited the flowers for nectar while *Pontia daplidice* and *Callerebia nirmala* were recorded

only once visiting the flowers. We also observed *Vanessa cardui* (Nymphalidae) utilizing *C. wallichii* as a larval host plant. During the study period, *C. wallichii* was the only species that attracted diverse butterfly species.

Cirsium has been studied in terms of nectar source by several workers such as Robertson (1928) who reported 14 species of Lepidoptera foraging on *C. vulgare*, eight species on *C. altissimum* and nine species each on *C. discolor* and *C. pumilum*. Thirty-three pollinators including 15 species of butterflies visiting *C. verutum* have been reported from the western Himalaya (Kapkoti et al. 2016). Although, it is used as a medicinal plant by the tribal people of the Himalaya (Uniyal et al. 2011), *C. wallichii* has never been reported as an important forage. The present communication highlights the importance of *C. wallichii* as a key nectar source for a large number of butterfly species though the plant is considered as a weed. The visits of several species of butterflies to *C. wallichii* could be attributed to the



Image 2. Butterfly species visiting *Cirsium wallichii*: A—*Celaenorrhinus dhanada* | B—*Seseria dohertyi* | C—*Lobocla liliana* | D—*Celaenorrhinus munda* | E—*Aeromachus stigmata* | F—*Pedesta masuriensis* | G—*Potanthus dara* | H—*Notocrypta feisthamelii* | I—*Polytremis discrete* | J—*Parnara* sp. | K—*Celastrina huegelii* | L—*Chilades pandava*. © Bitupan Boruah.



Image 3. Butterfly species visiting *Cirsium wallichii*. A—*Rapala manea* | B—*Lycaena phlaeas* | C—*Heliophorus sena* | D—*Spindasis nipalicus* | E—*Deudorix epijarbas* | F—*Lampides boeticus* | G—*Belenois aurota* | H—*Pontia daplidice* | I—*Gonepteryx rhamni* | J—*Pieris brassicae* | K—*Aporia leucodice* | L—*Colias fieldii*. © Bitupan Boruah.



Image 4. Butterfly species visiting *Cirsium wallichii*: A—*Colias erate* | B—*Aporia agathon caphusa* | C—*Aporia agathon agathon* | D—*Papilio bianor* | E—*Graphium agamemnon* | F—*Graphium sarpedon* | G—*Graphium cloanthus* | H—*Papilio protenor* | I—*Danaus genutia* | J—*Parantica aglea* | K—*Argynnis childreni* | L—*Lasiommata schakra*. © Bitupan Boruah.



Image 5. Butterfly species visiting *Cirsium wallichii*: A & B—*Vanessa cardui* | C—*Ypthima nareda* | D & E—*Argynnis hyperbius* | F—*Aglais caschmirensis* | G & H—*Vanessa indica* | I—*Callerebia annada caeca* | J—*Callerebia nirmala* | K & L—*Euploea mulciber*. © Bitupan Boruah.

hexose-rich sugar and strong amino acid content in the florets. This characteristic of the plants belonging to the family Asteraceae has been reported by Baker & Baker (1983). As observed on *Wendlandia tinctoria* (Raju et al. 2011), clustered flowering of *C. wallichii* also have benefited the butterflies thus, reducing searching time. Thistle in the Himalaya such as *C. verutum* has been found as an important forage (Kapkoti et al. 2016) and it proves to be an important resource for butterflies in the Benog Wildlife Sanctuary, Mussoorie. This study indicates that there is a need for further studies to understand the role of *C. wallichii* in sustaining butterfly diversity at landscape level during summer season.


References

- Baker, H.G. & I. Baker (1983). Floral nectar sugar constituents in relation to pollinator type, pp. 117–141. In: Jones, C.E. & R.J. Little (eds.). *Handbook of Experimental Pollination Biology*. Scientific and Academic Editions, New York, 558pp.
- Champion, H.G. & S.K. Seth (1968). *A Revised Survey of Forest Types of India*. Govt. of India Press, Delhi, 404pp.
- Courtney, S.P. (1986). The ecology of pierid butterflies: dynamics and interactions. *Advances in Ecological Research* 15: 51–131.
- Evans, W.H. (1932). *Identification of Indian Butterflies* 2nd edition. Bombay Natural History Society, Bombay, 464pp.
- Kapkoti, B., R.K. Joshi & R.S. Rawal (2016). Thistle (*Cirsium verutum*): An important forage for pollinators in Kumaun, West Himalaya. *National Academy Science Letters* 39(5): 395–399. <https://doi.org/10.1007/s40009-016-0501-x>
- Kehimkar, I. (2016). *Butterflies of India*. Bombay Natural History Society, Mumbai, 528pp.
- Kitahara, M., M. Yumoto & T. Kobayashi (2008). Relationship of butterfly diversity with nectar plant species richness in and around the Aokigahara primary woodland of Mount Fuji, Central Japan. *Biodiversity and Conservation* 17(11): 2713–2734. <https://doi.org/10.1007/s10531-007-9265-4>
- Kumar, A., M. Mitra, G. Singh & G.S. Rawat (2012). An inventory of the flora of Binog Wildlife Sanctuary, Mussoorie, Garhwal Himalaya. *Indian Journal of Fundamental and Applied Life Sciences* 2(1): 281–299.
- Kunte, K. (2000). *India – A Lifescope: Butterflies of Peninsular India*. Universities Press, Hyderabad, 254pp.
- Mabberley, J.D. (2008). *Mabberley's Plant Book: A Portable Dictionary of Plants, their Classification and Uses* 3rd edition. Cambridge University Press, UK, 1102pp.
- Mukherjee, S., S. Banerjee, P. Basu, G. Saha & G. Aditya (2015). *Lantana camara* and butterfly abundance in an urban landscape: benefits for conservation or species invasion? *Ekológia (Bratislava)* 34(4): 309–328. <https://doi.org/10.1515/eko-2015-0029>
- Murphy, D. & B.A. Wilcox (1986). Butterfly diversity in natural habitat fragments: a test of the validity of vertebrate-based management, pp. 287–292. In: Verner, J., M. Morrison & C.J. Ralph (eds.). *Wildlife 2000, Modelling Habitat Relationships of Terrestrial Vertebrates*. University of Wisconsin Press, Madison, 470pp.
- Nimbalkar, R.K., S.K. Chandekar & S.P. Khunte (2011). Butterfly diversity in relation to nectar food plants from Bhor Tahsil, Pune District, Maharashtra, India. *Journal of Threatened Taxa* 3(3): 1601–1609. <https://doi.org/10.11609/JoTT.o2612.1601-9>
- Raju, A.J.S., A. Bhattacharya & S.P. Rao (2004). Nectar host plants of some butterfly species at Visakhapatnam. *Scientific and Culture* 70 (4–5): 187–190.
- Raju, A.J.S., K.V. Ramana & P.V. Lakshmi (2011). *Wendlandia tinctoria* (Roxb.) DC. (Rubiaceae), a key nectar source for butterflies during the summer season in the southern Eastern Ghats, Andhra Pradesh, India. *Journal of Threatened Taxa* 3(3): 1594–1600. <https://doi.org/10.11609/JoTT.o2503.1594-600>
- Robertson, C. (1928). *Flowers and insects: Lists of Visitors of Four Hundred and Fifty-Three Flowers*. The Science Press Printing Company, Lancaster, PA, 221pp. <https://doi.org/10.5962/bhl.title.11538>
- Sahli, R., C. Rivière, C. Dufloer, C. Beaufay, C. Neut, J. Bero, T. Hennebelle, V. Roumy, R. Ksouri, J. Quetin-Leclercq & S. Sahpaz (2017). Antiproliferative and antibacterial activities of *Cirsium scabrum* from Tunisia. *Evidence-Based Complementary and Alternative Medicine* 2017: 1–9. <https://doi.org/10.1155/2017/7247016>
- Tooker, J.F., P.F. Reagel & L.M. Hanks (2002). Nectar sources of day-flying Lepidoptera of central Illinois. *Annals of Entomological Society of America* 95(1): 84–96.
- Uniyal, S.K., V. Sharma & P. Jamwal (2011). Folk medicinal practices in Kangra District of Himachal Pradesh, Western Himalaya. *Human Ecology* 39(4): 479–488. <https://doi.org/10.1007/s10745-011-9396-9>





www.threatenedtaxa.org



The Journal of Threatened Taxa (JoTT) is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at www.threatenedtaxa.org. All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use, reproduction, and distribution of articles in any medium by providing adequate credit to the author(s) and the source of publication.

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

October 2020 | Vol. 12 | No. 14 | Pages: 16927–17062

Date of Publication: 26 October 2020 (Online & Print)

DOI: 10.11609/jott.2020.12.14.16927-17062

Article

Elevational pattern and seasonality of avian diversity in Kaligandaki River Basin, central Himalaya

– Juna Neupane, Laxman Khanal, Basant Gyawali & Mukesh Kumar Chalise, Pp. 16927–16943

Communications

A highway to hell: a proposed, inessential, 6-lane highway (NH173) that threatens the forest and wildlife corridors of the Western Ghats, India

– H.S. Sathya Chandra Sagar & Mrunmayee, Pp. 16944–16953

Species diversity and feeding guilds of birds in Malaysian agarwood plantations

– Nor Nasibah Mohd Jamil, Husni Ibrahim, Haniza Hanim Mohd Zain & Nur Hidayat Che Musa, Pp. 16954–16961

Evaluating performance of four species distribution models using Blue-tailed Green Darner *Anax guttatus* (Insecta: Odonata) as model organism from the Gangetic riparian zone

– Kritish De, S. Zeeshan Ali, Niladri Dasgupta, Virendra Prasad Uniyal, Jeyaraj Antony Johnson & Syed Ainul Hussain, Pp. 16962–16970

Butterfly species richness and diversity in rural and urban areas of Sirajganj, Bangladesh

– Sheikh Muhammad Shaburul Imam, Amit Kumer Neogi, M. Ziaur Rahman & M. Sabbir Hasan, Pp. 16971–16978

Chroococcalean blue green algae from the paddy fields of Satara District, Maharashtra, India

– Sharada Jagannath Ghadage & Vaneeta Chandrashekhar Karande, Pp. 16979–16992

Short Communications

Avifaunal diversity along the riverine habitats of Papikonda National Park, Andhra Pradesh, India

– Paromita Ray, Giridhar Malla, Upma Manral, J.A. Johnson & K. Sivakumar, Pp. 16993–16999

Medetomidine may cause heart murmur in Cougars and Jaguars: case report

– Thiago Cavalheri Luczinski, Gediendson Ribeiro de Araújo, Matheus Folgareini Silveira, Murillo Dapará Kirnew, Roberto Andres Navarrete, Jorge Aparecido Salomão-Jr, Letícia Alecho Requena, Jairo Antonio Melo dos Santos, Marcell Hideki Koshiyama, Cristiane Schilbach Pizzutto & Pedro Nacib Jorge-Neto, Pp. 17000–17002

Description of a new species of *Omyomymar* Schauff from India with a key to Oriental species and first report of *Palaeoneura markhoddlei* Triapitsyn (Hymenoptera: Mymaridae) from the Indian subcontinent

– H. Sankararaman & S. Manickavasagam, Pp. 17003–17008

Incursion of the killer sponge *Terpios hoshinota* Rützler & Muzik, 1993 on the coral reefs of the Lakshadweep archipelago, Arabian Sea

– Rocktim Ramen Das, Chemmencheri Ramakrishnan Sreeraj, Gopi Mohan, Kottarathil Rajendran Abhilash, Vijay Kumar Deepak Samuel, Purvaja Ramchandran & Ramesh Ramchandran, Pp. 17009–17013

Contribution to the macromycetes of West Bengal, India: 63–68

– Rituparna Saha, Debal Ray, Anirban Roy & Krishnendu Acharya, Pp. 17014–17023

Notes

A rare camera trap record of the Hispid Hare *Caprolagus hispidus* from Dudhwa Tiger Reserve, Terai Arc Landscape, India

– Sankarshan Rastogi, Ram Kumar Raj & Brides Kumar Chauhan, Pp. 17024–17027

First distributional record of the Lesser Adjutant *Leptoptilos javanicus* Horsfield, 1821 (Ciconiiformes: Ciconiidae) from Sindhuli District, Nepal

– Badri Baral, Sudeep Bhandari, Saroj Koirala, Parashuram Bhandari, Ganesh Magar, Dipak Raj Basnet, Jeevan Rai & Hem Sagar Baral, Pp. 17028–17031

First record of African Sailfin Flying Fish *Parexocoetus mento* (Valenciennes, 1847) (Beloniformes: Exocoetidae), from the waters off Andaman Islands, India

– Y. Gladston, S.M. Ajina, J. Praveenraj, R. Kiruba-Sankar, K.K. Bineesh & S. Dam Roy, Pp. 17032–17035

A first distribution record of the Indian Peacock Softshell Turtle *Nilssonina hurum* (Gray, 1830) (Reptilia: Testudines: Trionychidae) from Mizoram, India

– Gospel Zothanmawia Hmar, Lalbiakzuala, Lalmuansanga, Dadina Zote, Vanlalhrauaia, Hmar Betlu Ramengmawii, Kulendra Chandra Das & Hmar Tlawmte Lalremsanga, Pp. 17036–17040

A frog that eats foam: predation on the nest of *Polypedates* sp. (Rhacophoridae) by *Euphylyctis* sp. (Dicroglossidae)

– Pranoy Kishore Borah, Avrajjal Ghosh, Bikash Sahoo & Aniruddha Datta-Roy, Pp. 17041–17044

New distribution record of two endemic plant species, *Euphorbia kadapensis* Sarojin. & R.R.V. Raju (Euphorbiaceae) and *Lepidagathis keralensis* Madhus. & N.P. Singh (Acanthaceae), for Karnataka, India

– P. Raja, N. Dhatchanamoorthy, S. Soosairaj & P. Jansirani, Pp. 17045–17048

Cirsium wallichii DC. (Asteraceae): a key nectar source of butterflies

– Bitupan Boruah, Amit Kumar & Abhijit Das, Pp. 17049–17056

Hyecoum pendulum L. (Papaveraceae: Ranunculales): a new record for the flora of Haryana, India

– Naina Palria, Nidhan Singh & Bhoo Dev Vashistha, Pp. 17057–17059

Addendum

Erratum and addenda to the article 'A history of primatology in India'

– Mewa Singh, Mridula Singh, Honnavalli N. Kumara, Dilip Chetry & Santanu Mahato, Pp. 17060–17062

Member



Publisher & Host

