OPEN ACCESS



All articles published in the Journal of Threatened Taxa are registered under Creative Commons Attribution 4.0 International License unless otherwise mentioned. JoTT allows unrestricted use of articles in any medium, reproduction and distribution by providing adequate credit to the authors and the source of publication.



Journal of Threatened Taxa

The international journal of conservation and taxonomy

www.threatenedtaxa.org

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

Note

A SIGHTING OF *PLASTINGIA NAGA* (DE NICÉVILLE, [1884]) (LEPIDOPTERA: HESPERIIDAE: HESPERIINAE) FROM CENTRAL ASSAM, INDIA

Gaurab Nandi Das, Arajush Payra & Bitupan Boruah

26 September 2016 | Vol. 8 | No. 11 | Pp. 9382–9383 10.11609/jott.1985.8.11.9382-9383



For Focus, Scope, Aims, Policies and Guidelines visit http://threatenedtaxa.org/About_JoTT.asp
For Article Submission Guidelines visit http://threatenedtaxa.org/Submission_Guidelines.asp
For Policies against Scientific Misconduct visit http://threatenedtaxa.org/JoTT_Policy_against_Scientific_Misconduct.asp
For reprints contact <info@threatenedtaxa.org>

Partner



Publisher/Host





ISSN 0974-7907 (Online) ISSN 0974-7893 (Print)

OPEN ACCESS



Plastingia naga (de Nicéville, [1884]) (Hesperiidae) is a widely distributed butterfly that occurs from Assam (northeastern India) east to Myanmar, Malaysia, Thailand, Laos, Indonesia, Borneo and Philippines (Evans 1932; 1949 & http://yutaka.it-n.jp/hes/92120001.html). This species is "rare" in northeastern India

(Evans 1932) but it is moderately common in the Malayan region. As per Butterflies of Singapore, it is a moderately common butterfly in Singapore which has been frequently sighted near the larval host plant Fishtail Palm *Caryota mitis*.

Plastingia naga was first described by de Nicéville (1884) as Hesperia naga from Sibsaghar (upper Assam), when one female specimen of this species was collected, which is preserved in the Indian Museum, Calcutta. Later, it was also collected by Moore & Swinhoe from the Khasi and Jaintia Hills, Meghalaya (Swinhoe 1912–1913).

On 10 Oct 2014, at around 10:00hr, a solitary individual of *Plastingia naga* was sighted in Panbari Reserve Forest, central Assam (Fig. 1). The individual was sitting on a leaf in the shadows at a height of about 1.5m from the ground. The habitat predominantly comprised evergreen trees and shrubs. The authors photographed the species (Image 1 a,b) after which it took wing and did not return. Regular visits to the same site yielded no further sightings of *Plastingia naga*.

Plastingia naga has a blackish-brown ground colour with distinct pale white spots on the underside. Its upper side is marked with obscure yellow spots, and a small spot in cell may be present. The female is

A SIGHTING OF *PLASTINGIA NAGA* (DE NICÉVILLE, [1884]) (LEPIDOPTERA: HESPERIIDAE: HESPERIINAE) FROM CENTRAL ASSAM, INDIA

Gaurab Nandi Das¹, Arajush Payra² & Bitupan Boruah³

Bokakhat Town, Golaghat, Assam 785612, India
 Ramnagar, Purba Medinipur, West Bengal 721441, India
 Durgapur, Kaziranga, Assam 785609, India
 gaurab68nandidas@gmail.com (corresponding author),
 arajushpayra@gmail.com, ³ bitupan.kaz@gmail.com

morphologically similar to the male. The wing span of the species is 33–38 mm (Evans 1932).

Panbari Reserve Forest lies between Kaziranga National Park and the foothills of West Karbi Hills in central Assam. The reserve is bounded by National Highway 37 on the north, a human settlement on the east, and a tea garden and Karbi foot hills on the southwest. The altitude of the area varies from 80–360 m. The butterflies of this area were well studied by Gogoi (2013, 2015).

In 2013, Gogoi reported 137 species of the Hesperiidae butterfly from Panbari Reserve Forest and recorded three species of Lancers; Salanoemia fuscicornis, Pyroneura margherita margherita, Pyroneura niasana burmana but did not report any Plastingia naga (Gogoi 2013).

There are only a few recent records of *Plastingia naga* from mainland India. In recent times it has been sighted from the Siju Wildlife Sanctuary, South Garo Hills, on 22 May 2015 (Karmakar 2016). Karmakar reared *Plastingia naga* from its larval stage. This species has also been sighted from the Barail Range in September 2015

DOI: http://dx.doi.org/10.11609/jott.1985.8.11.9382-9383 | ZooBank: urn:lsid:zoobank.org:pub:172C10E9-03D8-406A-AA82-4419B777B5A5

Editor: Sanjay Sondhi, Titli Trust, Dehradun, India.

Date of publication: 26 September 2016 (online & print)

Manuscript details: Ms # 1985 | Received 01 April 2016 | Final received 17 May 2016 | Finally accepted 02 September 2016

Citation: Das, G.N., A. Payra & B. Boruah (2016). A sighting of *Plastingia naga* (de Nicéville, [1884]) (Lepidoptera: Hesperiidae: Hesperiinae) from central Assam, India. *Journal of Threatened Taxa* 8(11): 9382–9383; http://dx.doi.org/10.11609/jott.1985.8.11.9382-9383

Copyright: © Das et al. 2016. Creative Commons Attribution 4.0 International License. JoTT allows unrestricted use of this article in any medium, reproduction and distribution by providing adequate credit to the authors and the source of publication.

Funding: None.

 $\label{lem:conflict} \textbf{Conflict of Interest:} \ \textbf{The authors declare no competing interests.}$

Acknowledgements: We would to like thank to Mr. Isaac kehimkar for his help in the preparing the manuscript and also like to thank to Monsoon Jyoti Gogoi for his constant support.





Image 1 (a & b). Underside of Plastingia naga at Panbari Reserve Forest, central Assam.



Figure 1. Location of Panbari Reserve Forest, central Assam.

(Monsoon Jyoti Gogoi pers. comm. 2015). This record from Panbari shows the importance of these habitats for rare species in the last remaining forest patches of central Assam and the need to conserve these habitats. Further, long term monitoring is necessary to reveal the ecology and distribution of such butterfly species in northeastern India.

References

Anonymous (2014). Plastingia naga de Nicèville, 1884. Butterflies of Singapore. http://butterflycircle.blogspot.in/2013/07/lifehistoryofchequeredlancer.html. (Accessed on October 2014).

de Nicéville (1884). On new and little known Rhopalocera from the Indian Region. *Journal of Asiatic Society, Bengal* 52 Pt.II (2/4): 65–91, pl. 1, 9–10.

Evans, W.H. (1932). *The Identification of Indian Butterflies - 2nd Edition.* Bombay Natural History Society, Mumbai, India, 464pp+32pls.

Evans, W.H. (1949). A Catalogue of the Hesperiidae of Europe, Asia and Australia in the British Museum (Natural History). Trustees of the British Museum, London, 502pp.

Gogoi, M.J. (2013). Notes on some skipper butterflies (Lepidoptera: Hesperiidae) from Panbari Forest and its adjoining areas, Kaziranga-Karbi Anglong, upper Assam, India. *Journal of Threatened Taxa* 5(13): 4759–4768; http://dx.doi.org/10.11609/JoTT.o3340.4759-68

Gogoi, M.J. (2015). Observations on lycaenid butterflies from Panbari Reserve Forest and adjoining areas, Kaziranga, Assam, northeastern India. *Journal of Threatened Taxa* 7(15): 8259–8271; http://dx.doi.org/10.11609/jott.2467.7.15.8259-8171

Karmakar, T. (2016). Plastingia naga de Nicéville, 1883 - Silver-spotted Lancer. In: Kunte, K., P. Roy, S. Kalesh & U. Kodandaramaiah (eds.). Butterflies of India, v. 2.23. Indian Foundation for Butterflies. http:// www.ifoundbutterflies.org/sp/2950/Plastingia-naga.

Swinhoe, C. (1912–1913). Lepidoptera Indica. Part X. Rhopalocera-Hesperiidae. Lovell, Reeve & Co. Ltd., London, 364pp+757–835pls.







All articles published in the Journal of Threatened Taxa are registered under Creative Commons Attribution 4.0 International License unless otherwise mentioned. JoTT allows unrestricted use of articles in any medium, reproduction and distribution by providing adequate credit to the authors and the source of publication.

ISSN 0974-7907 (Online); ISSN 0974-7893 (Print)

September 2016 | Vol. 8 | No. 11 | Pages: 9289–9396 Date of Publication: 26 September 2016 (Online & Print) DOI: 10.11609/jott.2016.8.11.9289-9396

www.threatenedtaxa.org

Review

Distribution records and extended range of the Sri Lanka Frogmouth Batrachostomus moniliger (Aves: Caprimulgiformes: Podargidae) in the Western Ghats: a review from 1862 to 2015

-- Anil Mahabal, Sanjay Thakur & Rajgopal Patil, Pp. 9289-9305

Short Communications

Small carnivores of Parambikulam Tiger Reserve, southern Western Ghats, India

-- R. Sreehari & P.O. Nameer, Pp. 9306-9315

First record of the Diadem Leaf-Nosed Bat *Hipposideros diadema* (E. Geoffroy, 1813) (Chiroptera: Hipposideridae) from the Andaman Islands, India with the possible occurrence of a hitherto unreported subspecies

-- Bhargavi Srinivasulu, Aditya Srinivasulu, Chelmala Srinivasulu, Tauseef Hamid Dar, Asad Gopi & Gareth Jones, Pp. 9316–9321

New distribution records of *Mesoclemmys vanderhaegei* (Testudines: Chelidae) from southeastern Brazil, including observations on reproduction

-- Fábio Maffei, Bruno Tayar Marinho do Nascimento, Guilherme Marson Moya & Reginaldo José Donatelli, Pp. 9322–9326

Spiders (Arachnida: Araneae) of Gujarat University Campus, Ahmedabad, India with additional description of *Eilica tikaderi* (Platnick, 1976)

-- Dhruv A. Prajapati, Krunal R. Patel, Sandeep B. Munjpara, Shiva S. Chettiar & Devendrasinh D. Jhala, Pp. 9327–9333

New records of Termite species from Kerala (Isoptera: Termitidae)

-- Poovoli Amina, K. Rajmohana, K.V. Bhavana & P.P. Rabeeha, Pp. 9334–9338

Odonata (Insecta) diversity of southern Gujarat, India

-- Darshana M. Rathod, B.M. Parasharya & S.S. Talmale, Pp. 9339–9349

An update on the distribution pattern and endemicity of three lesser-known tree species in the Western Ghats, India

-- K. Sankara Rao, N.V. Page, A.N. Sringeswara, R. Arun Singh & Imran Baig, Pp. 9350–9355

Heavy metal distribution in mangrove sediment cores from selected sites along western coast of India

-- P. Vidya & Rajashekhar K. Patil, Pp. 9356–9364

Notes

New distribution record of the Bhutan Takin *Budorcas taxicolor* whitei Hodgson, 1850 (Cetartiodactyla: Bovidae) in Bhutan -- Tashi Dhendup, Tshering Tempa, Tsethup Tshering & Nawang Norbu, Pp. 9365–9366

Recent records and distribution of the Indian Brown Mongoose Herpestes fuscus Gray, 1837 (Mammalia: Carnivora: Herpestidae) from the southern Western Ghats, India

-- R. Sreehari, Sandeep Das, M. Gnanakumar, K.P. Rajkumar, K.A. Sreejith, Navaneeth Kishor, Dhaneesh Bhaskar, P.S. Easa & P.O. Nameer, Pp. 9367–9370

First record of Dobson's Long-tongued Fruit Bat *Eonycteris spelaea* (Dobson, 1871) (Mammalia: Chiroptera: Pteropodidae) from Kerala, India

-- P.O. Nameer, R. Ashmi, Sachin K. Aravind & R. Sreehari, Pp. 9371–9374

Road kills of the endemic snake Perrotet's Shieldtail *Plectrurus perrotetii,* Dumeril, 1851 (Reptilia: Squamata: Uropeltidae) in Nilgiris, Tamil Nadu, India

-- P. Santhoshkumar, P. Kannan, B. Ramakrishnan, A. Veeramani, A. Samson, S. Karthick, J. Leonaprincy, B. Nisha, N. Dineshkumar, A. Abinesh, U. Vigneshkumar & P. Girikaran, Pp. 9375–9376

Reappearance of the rare Shingle Urchin Colobocentrotus (Podophora) atratus (Camarodonta: Echinometridae) after eight decades from the rocky shore of Kodiyaghat (Port Blair), South Andaman Islands, India

-- Vikas Pandey & T. Ganesh, Pp. 9377–9380

Sallywalkerana, a replacement name for Walkerana Dahanukar et al. 2016 (Anura: Ranixalidae)

-- Neelesh Dahanukar, Nikhil Modak, Keerthi Krutha, P.O. Nameer, Anand D. Padhye & Sanjay Molur, P. 9381

A sighting of *Plastingia naga* (de Nicéville, [1884]) (Lepidoptera: Hesperiidae: Hesperiinae) from central Assam, India

-- Gaurab Nandi Das, Arajush Payra & Bitupan Boruah, Pp. 9382–9383

A note on the taxonomy, field status and threats to three endemic species of *Syzygium* (Myrtaceae) from the southern Western Ghats,

-- R. Ramasubbu, C. Divya & S. Anjana, Pp. 9384–9390

Arnebia nandadeviensis Sekar & Rawal (Boraginaceae) a new synonym of Onosma bracteata Wall.

-- Umeshkumar L. Tiwari, Pp. 9391–9393

Exosporium monanthotaxis Piroz. (Fungi: Ascomycota: Pezizomycotina) from Biligirirangan Hills, southern India

-- Rashmi Dubey & Shreya Sengupta, Pp. 9394–9396



