

OPEN ACCESS



The Journal of Threatened Taxa is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at www.threatenedtaxa.org. All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use of articles in any medium, reproduction, and distribution by providing adequate credit to the authors and the source of publication.

Journal of Threatened Taxa

Building evidence for conservation globally

www.threatenedtaxa.org

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

SHORT COMMUNICATION

THE GILLED MUSHROOM *AMANITA SPISSACEA* (AMANITACEAE): A NEW REPORT FOR INDIA

Hmar Lalrinawmi, John Zothanzama, Benjamin W. Held,
Josiah M.C. Vabeikhokey, Zohmangaiha & Robert A. Blanchette

26 September 2018 | Vol. 10 | No. 10 | Pages: 12413–12417
10.11609/jott.3471.10.10.12413-12417



For Focus, Scope, Aims, Policies and Guidelines visit <http://threatenedtaxa.org/index.php/JoTT/about/editorialPolicies#custom-0>
For Article Submission Guidelines visit <http://threatenedtaxa.org/index.php/JoTT/about/submissions#onlineSubmissions>
For Policies against Scientific Misconduct visit <http://threatenedtaxa.org/index.php/JoTT/about/editorialPolicies#custom-2>
For reprints contact info@threatenedtaxa.org

Partners



محمود بن زايد
للمحافظة على
الكائنات الحية
The Mohamed bin Zayed
SPECIES CONSERVATION FUND



Member



Publisher & Host





THE GILLED MUSHROOM *AMANITA SPISSACEA* (AMANITACEAE): A NEW REPORT FOR INDIA

Hmar Lalrinawmi¹ , John Zothanzama² , Benjamin W. Held³ , Josiah M.C. Vabeikhochei⁴ ,
Zohmangaiha⁵  & Robert A. Blanchette⁶ 

ISSN 0974-7907 (Online)
ISSN 0974-7893 (Print)

^{1,2,4,5} Department of Environmental Sciences, Mizoram University, Tanhril, Mizoram 796004, India

^{3,6} University of Minnesota, 495 Borlaug Hall, 1991 Upper Buford Circle, St. Paul, MN 55108, United States

¹ hmarlalrinawmi@gmail.com, ² john_zza@yahoo.co.in (corresponding author), ³ bhheld@umn.edu,

⁴ mcjosiahmathipy@gmail.com, ⁵ xohmaa1990@gmail.com, ⁶ robertb@umn.edu

OPEN ACCESS



Abstract: Mizoram is regarded as one of the biodiversity hotspots of the World owing to the diverse group of flora and fauna documented here. Information regarding the macrofungi, however, is very limited. For this reason, a systematic study of mushrooms from Mizoram was undertaken and during the field survey, *Amanita spissacea* was collected and identified. This is the first report of this mushroom from India. This species was identified on the basis of its morphological and microscopic characteristics as well as molecular characterization of the ITS region of rDNA. Phylogenetic analysis also confirmed that *A. spissacea* was a distinct species from *A. fritillaria*, *A. sepiacea*, *A. citrina* and other closely related species *Amanita* section *Valideae*.

Keywords: Macrofungi, Mizoram, phylogeny, taxonomy.

Mizoram lies in northeastern India sharing its borders with Assam, Manipur and Tripura and has international borders with Bangladesh and Myanmar. It covers a geographical area of 21,081 km² and lies between 21.966–24.583°N and 91.250–92.483°E. The Tropic of Cancer passes through the state at 23.500°N (Mizoram Remote Sensing Application Centre 2009).

Amanita Pers., is a well known mushroom genus with global distribution comprising both edible and poisonous species which are usually mycorrhizal

symbionts with plants. The genus *Amanita* Pers., contains about 500 species worldwide (Kirk et al. 2008), and for some time, only 66 species were reported from India (Bhatt et al. 2003; Semwal et al. 2005, 2007; Vrinda et al. 2005). Recently, a number of reports have been added to the list from several researchers (Singh & Kaur 2016; Bhatt et al. 2017) with the latest report of 80 species of Amanitaceae being listed including 73 species of *Amanita* reported from different parts of India (Verma & Pandro 2018).

During the course of macro-fungal foray to different parts of Mizoram, *Amanita spissacea* S. Imai was collected and identified. This species is described and illustrated for the first time from India.

MATERIALS AND METHODS

Study Area

Collections of mushrooms growing on soil was done at Mizoram University Campus which is located in the Western side at a distance of about 15 km away from the state capital, Aizawl, just below Tanhril Village. The Mizoram University Campus is about 980 acres in area

DOI: <https://doi.org/10.11609/jott.3471.10.10.12413-12417>

Editor: R.K. Verma, Tropical Forest Research Institute, Jabalpur, India.

Date of publication: 26 September 2018 (online & print)

Manuscript details: Ms # 3471 | Received 28 April 2017 | Final received 02 August 2018 | Finally accepted 23 August 2018

Citation: Lalrinawmi, H., John Zothanzama, Benjamin Held, Josiah M.C. Vabeikhochei, Zohmangaiha & Robert A. Blanchette (2018). The gilled mushroom *Amanita spissacea* (Amanitaceae): a new report for India. *Journal of Threatened Taxa* 10(10): 12413–12417; <https://doi.org/10.11609/jott.3471.10.10.12413-12417>

Copyright: © Lalrinawmi et al. 2018. Creative Commons Attribution 4.0 International License. JoTT allows unrestricted use of this article in any medium, reproduction and distribution by providing adequate credit to the authors and the source of publication.

Funding: UGC-MZU (Non-NET) Fellowship.

Competing interests: The authors declare no competing interests.

Acknowledgements: The authors thank Dr. Karlyn Eckman and collaborators involved in the joint cooperative research projects between Mizoram University, India and the University of Minnesota, USA. The authors also thank Dr. Surajit De Mandal, Research Associate, Department of Biotechnology, Mizoram University.



and lies between 23.756–23.726°N & 92.644–92.673°E. The elevation ranges from 330–880 m.

Morphological study

Macro-morphological descriptions were based on field notes and color photographs of the macrofungi. Micro-morphological data was obtained from the dried specimens with the aid of a light microscope after sectioning and staining with cotton-blue. Spore prints were taken by placing the fresh specimen on a microslide. Descriptions of spore shapes are based on the study reported by Bas (1969).

Phylogenic study

DNA isolation, amplification and sequencing:

Molecular methods were performed following Zothanzama et al. (2016), where DNA was extracted using a CTAB method, followed by amplification of the internal transcribed spacer region (ITS) of the rDNA and sequenced with both primers (ITS1F and ITS4B).

PCR amplification: PCR reactions were setup in 0.2ml centrifuge tubes that contained 12.5µl GoTaq Green Mastermix (Promega, Madison, WI), 9.5µl nuclease free water, 0.5µl bovine serum albumin (BSA), 1µl forward primer (5µM), 1µl reverse primer (5µM) and 1µl of fungal DNA template for a total reaction volume of 25.5µl. PCR was performed using primers ITS1-F (5'-CTT GGT CAT TTA GAG GAA GTA A-3') ITS4-B (5'-CAG GAG ACT TGT ACA CGG TCC AG-3') (White et al. 1990) with the following parameters; 94C for 5 minutes, followed by 35 cycles of 94C for 1 minute, 52C for 1 minute and 72C for 1 minute with a final extension step of 72C. PCR amplicons were verified by electrophoresis on a 1% agarose gel with SYBR green and visualized on a Gel Documentation System. Sequencing was performed using both primers by using Sanger sequencing using a ABI 3730xl DNA sequencer. Consensus sequences for contigs were trimmed and aligned using Bioedit sequence alignment editor. Sequences were then compared to those in GenBank database using the BLASTn (Altschul et al. 1990) search tool for similarities and submitted to Genbank.

Phylogenetic analysis: The ITS dataset was aligned with the MAFFT v7.222 (Katoh et al. 2002) and jModelTest 2.1.10 (Darriba et al. 2012) was used to determine the appropriate model for Bayesian analysis (HKY85). Phylogenetic analysis inferred from ITS sequences was performed using MrBayes 3.2.6 (Huelsenbeck & Ronquist 2001). 1.1×10^6 MCMC generations were used with a sampling frequency every 200 generations and the first 10% of sampled trees were discarded as burn-in.

RESULTS

Amanita spissacea S. Imai (Fig. 1 & Image 1)

Specimens examined: EVS/SF/0012, 27.v.2014, India, Mizoram, Aizawl, Mizoram University Campus and EVS/SF/0165, 01.vi.2016 (Image 2).

Basidiomata: Small to medium. 4–9 cm in diam., convex to plano-convex, grayish-brown in color, volva remnants on pileus as scattered felted to crust like patches, margin non-striated, non-appendeculate, incurved. Lamellae-sometimes forked, lamellulae- of several length. Stipe: 8–14 cm long, 0.5–1 cm wide, tapering toward apex, stuffed, white to grayish-brown with brown scales. Annulus membranous, grayish brown, apical. Bulbous base upto 1.5cm long, 1–3 cm thick, glabrous with dark brown spots. The upper part of the bulbous base of the stipe is covered with dark grey volval remnants in 2–5 dotted rings. Context - white and thin. Sporeprint: White. Spores: 7-9.8 x 6.8 - 8.5µm[Q=1.02,1.15]and are globose to subglobose, sometimes rarely broadly ellipsoid, amyloid, colourless, hyaline, thin walled and smooth. Basidia: Clavate, 35–45




Figure 1. A–D *Amanita spissacea*.
A - Fruiting body, B - Spores, C - Basidium, D - Marginal cell or lamellae edge cell (scale: A - 3cm, B - 8µm, C & D - 6.5µm)



Image 1. A–C - Fruiting body of *Amanita spissacea* in their natural habitat; D - Fruiting body of *Amanita spissacea* in laboratory (scale A&B - 2cm; C&D - 4cm).

× 8–11 µm, four spored, sterigmata 3.2–4.6 × 0.8–1.8 µm. Clamp connection absent. Lamellae edge cell: clavate, 35–45 × 7.5–9 µm.

Habitat: Solitary to scattered or gregarious on ground in a broad-leaved sub-tropical forest.

Molecular Phylogenetic analysis

The molecular phylogenetic analysis shown in figure (Fig. 2) involved 17 nucleotide sequences. The tree with the highest log likelihood (-2938.15) is shown. In the phylogenetic analysis, the specimen of *Amanita spissacea* from Mizoram (MZ10-KY940266, MZJR1-MG706138) is indicated in bold and clustered with *Amanita spissacea* from Belgium (KY747469), Republic of Korea (KM052550, KM052546) and Japan (AB015683).

DISCUSSION

In this study, we identified this species based on morphological, microscopic and molecular characteristics. This is the first report of *Amanita spissacea* from India. Results from sectioning of the fruiting body and observations of basidiospores indicated this *Amanita* species was most closely related to *A. spissacea*. Species identification based on morphological characteristics is difficult to differentiate from other closely related species such as *A. fritillaria*, *A. sepiacea*, *A. citrina* and others.

The present species has been reported and described for the first time by Imai (1933) and Gilbert (1940) as *Amplariella spissacea*. The macro and microscopic features of the present species well matched with the description given by Imai (1933) who described *Amanita*




Figure 2. Phylogenetic tree of *Amanita spissacea* collected in Mizoram (MZ 10-KY940266 & MZ JZR1-MG706138) and other closely related *Amanita* species. The tree is drawn to scale, with branch lengths measured in the number of substitutions per site.

spissacea as pileus with 6–10 cm, convex, then extended, dark chestnut, warted, white flesh, stalk 10–15 cm long, firm, bulbous base, covered with dark margin, scaly, membranous ring. Spores globose, 7–8 μ m, hyaline, apiculate.

Amanita spissacea is also closely related to *Amanita fritillaria* (Yang et al. 2001) and *Amanita sepiacea* (Imai, 1933). The macro and microscopic details are mostly identical but both *Amanita fritillaria* and *Amanita sepiacea* have spores broadly ellipsoid to ellipsoid, occasionally subglobose or ellipsoid, rarely globose and the upper part of the bulbous base of the stipe of *Amanita fritillaria* covered with dark grey volval remnants is only 2–4 rings while the former is 2–5 dotted rings. Moreover, the macroscopic feature of *Amanita sepiacea* is bigger in size as compared to *Amanita spissacea* with cap 6–15 cm diam., stipe 10–18 cm long, 1–2.5 cm thick and basal bulb 1.5–5.0 cm.



Image 2. Herbarium image of *Amanita spissacea*

Sequencing of the ITS region of rRNA and phylogenetic analysis further showed that the Mizoram sample matched GenBank accession *Amanita spissacea* from Belgium (KY747469), Republic of Korea (KM052550, KM052546) and Japan (AB015683) in a well-supported clade with *A. fritillaria* forming a sister clade. These results hence confirmed that the specimen of *Amanita* from Mizoram (MZ10-KY940266, MZJZR1-MG706138) is *Amanita spissacea*, a distinct species and separate from *A. fritillaria*, *A. sepiacea*, *A. citrina* and other previous reported *Amanita* species.

Mizoram is one of the northeastern states of India which is rich in mushroom flora. Like many other *Amanita* species, *A. spissacea* has been reported to be poisonous in China (Zhishu et al. 1993) and recent mushroom poisonings in Mizoram State (Zothanzama & Lalrinawmi 2015) are prompting efforts to identify mushrooms in this region that are poisonous. This report identifies this poisonous mushroom in India and confirms that it is a distinct species from other *Amanita* species. Limited information is available concerning the wild mushrooms found in Mizoram and further studies are needed to assess and document the wide variety of wild mushrooms that can be found in this region.

REFERENCES

- Altschul, S.F., W. Gish, W. Miller, E.W. Myers & D.J. Lipman (1990). Basic local alignment search tool. *Journal of Molecular Biology* 215(3): 403–410; [https://doi.org/10.1016/S0022-2836\(05\)80360-2](https://doi.org/10.1016/S0022-2836(05)80360-2)
- Bas, C. (1969). Morphology and subdivision of *Amanita* and a monograph of its section *Lepidella*. *Persoonia* 5(3): 285–579.
- Bhatt, R.P., R.E. Tulloss, K.C. Semwal, V.K. Bhatt, J.M. Moncalvo & S.L. Stephenson (2003). Amanitaceae reported from India. A critically annotated checklist. *Mycotaxon* 88: 249–270.
- Bhatt, R.P., T. Mehmood, P. Uniyal & U. Singh (2017). Six new records of genus *Amanita* (Amanitaceae) from Uttarakhand, India. *Current Research in Environmental & Applied Mycology* 7(3): 161–182
- Darriba, D., G.L. Taboada, R. Doallo & D. Posada (2012). jModelTest 2: more models, new heuristics and parallel computing. *Nat Methods* 9: 772–772; <https://doi.org/10.1038/nmeth.2109>
- Gilbert, E.J. (1940). Iconographia mycologica, Amanitaceae. *Iconographia Mycologica* 27: 1–198.
- Huelsenbeck J.P. & F. Ronquist (2001). MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics* 17: 754–755; <https://doi.org/10.1093/bioinformatics/17.8.754>
- Imai, S. (1933). Studies on the Agaricaceae of Japan I. Volvate Agarics in Hokkaido. *Botanical Magazine* (Tokyo) 47: 423–432.
- Katoh, K., K. Misawa, K. Kuma & T. Miyata (2002). MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Research* 30: 3059–3066; <https://doi.org/10.1093/nar/gkf436>
- Kirk, P.M., P.F. Cannon, D.W. Minter & J.A. Stalpers (2008). *Ainsworth's Dictionary of Fungi*. 10th Edition. CABI, Wallingford, UK.
- Mizoram Remote Sensing Application Centre (2009). Natural Resources Atlas of Mizoram. *MIRSAC Newsletter* 2(1): 1–4 < http://mirsac.nic.in/images/MIRSAC_News_July2010.pdf>
- Semwal, K.C., R.P. Bhatt & R.C. Upadhyay (2005). The genus *Amanita* from Garhwal Himalayas region of India. *Mushroom Research* 14 (2): 50–55.
- Semwal, K.C., R.E. Tulloss, R.P. Bhatt, S.L. Stephenson & R.C. Upadhyay (2007). New records of *Amanita* from Garhwal Himalaya, India-*Amanita* section *Amanita*. *Mycotaxon* 101: 331–348.
- Singh, Y. & M. Kaur (2016). Two species of genus *Amanita* from India. *World Journal of Pharmacy and Pharmaceutical Sciences* 5(5): 1054–1062.
- Verma, R.K. & V. Pandro (2018). Diversity and distribution of amanitaceous mushrooms in India, two new reports from Sal forest of central India. *Indian Journal of Tropical Biodiversity* 26(1): 42–54.
- Vrinda, K.B., C.K. Pradeep & S.S. Kumar (2005). Occurrence of a lesser known edible *Amanita* in the Western Ghats of Kerala. *Mushroom Research* 14(1): 5–8.
- White, T.J., T. Bruns, S. Lee & J.W. Taylor (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics, pp. 315–322. In: Innis, M.A., D.H. Gelfand, H.H. Sninsky & T.J. White (eds.). *PCR Protocols: A Guide to Methods and Applications*. New York Academic Press Inc.
- Yang, Z.L., T.H. Li & X.L. Wu (2001). Revision of *Amanita* collections made from Hainan, Southern China. *Fungal Diversity* 6: 149–165.
- Zhishu, B., Z. Guoyang & L. Taihui (1993). *The Macrofungus Flora of China's Guangdong Province*. The Chinese University Press, Hong Kong.
- Zothanzama, J. & H. Lalrinawmi (2015). Wild edible mushrooms of Mizoram: an overview of knowledge and potential, pp. 171–181. In: Eckman, K. & L. Ralte (eds.). *Integrated Land Use Management in the Eastern Himalayas Vol - 1*. Akansha Publishing House. New Delhi.
- Zothanzama, J., R.A. Blanchette, S. Redford, B. Held, Zohmangaiha & J.M.C. Vabeikhookhei (2016). Using molecular characterization as a tool for identification of fungi from Mizoram, India, pp. 165–177. In: Eckman, K. & L. Ralte (eds.). *Integrated Land Use Management in the Eastern Himalayas. Vol - 2*. Akansha Publishing House. New Delhi.





OPEN ACCESS



The Journal of Threatened Taxa is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at www.threatenedtaxa.org. All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use of articles in any medium, reproduction, and distribution by providing adequate credit to the authors and the source of publication.

ISSN 0974-7907 (Online); ISSN 0974-7893 (Print)

September 2018 | Vol. 10 | No. 10 | Pages: 12299–12442

Date of Publication: 26 September 2018 (Online & Print)

DOI: 10.11609/jott.2018.10.10.12299-12442

www.threatenedtaxa.org

Communications

Urban biodiversity: an insight into the terrestrial vertebrate diversity of Guwahati, India

-- Jayaditya Purkayastha, Pp. 12299–12316

Status of raptors in the Moyar River Valley, Western Ghats, India

-- N.R. Anoop, S. Babu, S. Bharathidasan & R. Nagarajan, Pp. 12317–12327

Species composition and abundance estimates of reptiles in selected agroecosystems in southern Western Ghats, India

-- Abhirami Mini Jayakumar & Paingamadathil Ommer Nameer, Pp. 12328–12336

Comparison of beach profiles conducive for turtle nesting in Andaman

-- Subramanian Narayani, Sasidharan Venu & Andrea Joan D'Silva, Pp. 12337–12343

Short Communications

A new record of the rare Hardwicke's Woolly Bat *Kerivoula hardwickii* (Horsfield, 1824) (Mammalia: Chiroptera: Vespertilionidae) after 23 years from a lowland rainforest of Sri Lanka

-- Dinesh Gabadage, Gayan Edirisinghe, Madhava Botejue, Kalika Perera, Thilina Surasinghe & Suranjan Karunaratna, Pp. 12344–12349

Alarming population status of the Grizzled Giant Squirrel *Ratufa macroura* (Mammalia: Rodentia: Sciuridae) in Chinna Wildlife Sanctuary, the Western Ghats, India

-- Kiran Thomas & Paingamadathil Ommer Nameer, Pp. 12350–12356

Distribution and population status of Sambar *Rusa unicolor* (Mammalia: Cetartiodactyla: Cervidae) from Aravalli landscape with a note on its first record from Aravalli Hills of Haryana, India

-- Paridhi Jain, Anchal Bhasin, Gautam Talukdar & Bilal Habib, Pp. 12357–12362

Delayed peracute capture myopathy in a Himalayan Ibex *Capra sibirica* (Mammalia: Cetartiodactyla: Bovidae)

-- Umar Nazir Zahid, Latief Mohammad Dar, Umar Amin, Showkat Ahmad Shah, Rashid Yahya Naqash, Dil Mohammed Makhdooni, Shayuaib Ahmad Kamil & Intesar Suhail, Pp. 12363–12367

Checklist of the avifauna of Sagareshwar Wildlife Sanctuary, Maharashtra, India

-- Sharad Datt Apte, Vijay Bhagwan Tuljapurkar & Girish Avinash Jathar, Pp. 12368–12375

The rediscovery of Rurk's Cat Skink *Ristella rurkii* Gray, 1839 (Reptilia: Ristellidae) with remarks on distribution and natural history

-- Sumaithangi Rajagopalan Ganesh, Pp. 12376–12381

Dietary assessment of five species of anuran tadpoles from northern Odisha, India

-- Syed Asrafuzzaman, Susmita Mahapatra, Jasmin Rout & Gunanidhi Sahoo, Pp. 12382–12388

Partners



المؤسسة محمد بن زايد
للمحافظة على
الكائنات الحية
The Mohamed bin Zayed
Species Conservation Fund



Member



Publisher & Host

