


OPEN ACCESS



The Journal of Threatened Taxa is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at www.threatenedtaxa.org. All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use of articles in any medium, reproduction, and distribution by providing adequate credit to the authors and the source of publication.



Journal of Threatened Taxa

Building evidence for conservation globally

www.threatenedtaxa.org

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

RESPONSE & REPLY

BEST PROTOCOLS FOR CYCAD PROPAGATION REQUIRE MORE RESEARCH

Thomas Edward Marler & Gil Naputi Cruz

26 September 2017 | Vol. 9 | No. 9 | Pp. 10738–10740
10.11609/jott.3812.9.9.10738-10740



For Focus, Scope, Aims, Policies and Guidelines visit http://threatenedtaxa.org/About_JoTT
For Article Submission Guidelines visit http://threatenedtaxa.org/Submission_Guidelines
For Policies against Scientific Misconduct visit http://threatenedtaxa.org/JoTT_Policy_against_Scientific_Misconduct
For reprints contact <info@threatenedtaxa.org>

Partner



Publisher/Host





ISSN 0974-7907 (Online)
ISSN 0974-7893 (Print)

OPEN ACCESS



RESPONSE TO ARTICLE ON ADVENTITIOUS ROOTING OF MATURE *CYCAS MICRONESICA*

S. Suresh Ramanan

WCCB (Wildlife Crime Control Bureau)
Volunteer, Sher-e-Kashmir University of
Agricultural Sciences and Technology of
Jammu, Jammu and Kashmir 180009, India
sureshramanan01@gmail.com

A recent article by Marler & Cruz (2017) in JoTT states that the stem cutting can be used as planting material to save the endangered cycads. I list a few questions and comments that may be addressed by the authors to clarify some points.

1. Generally, in vegetative propagation methodology, the new vegetative cutting are maintained in mist chambers to enhance the rooting percentage. Was there any methodology adopted in this study?

2. From the article it can be presumed that cuttings were maintained in open air nurseries. A mention of the climatic condition of the study site will help in understanding the success of vegetative propagation.

3. What was the criteria in choosing the concentration of indole-3-butyric acid (IBA), is there any previous study in this regard? Instead of comparing the efficiency of cycad specialist and a general forestry nursery specialist, the authors would have attempted to compare the different concentration of rooting hormone on root formation.

4. Further, a photo showing the root formation in stem cutting would have made the article more vibrant.

5. Regarding the success rate between the two nursery conditions, one having 34 and the other having 15 stem cuttings, what was the actual number of plants

that rooted out?

6. Has this sort of adventitious rooting been reported for other cycads?

These are important issues that an article propounding vegetative propagation of an endangered species requires clarification.

Reference

Marler, T.E. & G.N. Cruz (2017). Adventitious rooting of mature *Cycas micronesica* K.D. Hill (Cycadales: Cycadaceae) tree stems reveals moderate success for salvage of an endangered cycad. *Journal of Threatened Taxa* 9(4): 10565–10570; <http://doi.org/10.11609/jott.3523.9.8.10565-10570>



BEST PROTOCOLS FOR CYCAD PROPAGATION REQUIRE MORE RESEARCH

Thomas Edward Marler¹ & Gil Naputi Cruz²

^{1,2} Western Pacific Tropical Research Center, UOG Station,
University of Guam, Mangilao, Guam 96923, USA

¹thomas.marler@gmail.com (corresponding author),

²gilcruz@triton.uog.edu

Knowledge of cycad biology and conservation is limited by the paucity of empirical studies. When adaptive management research protocols are envisioned in conservation programs, the best available science is first amassed, but then is distilled down to prioritize study methods. The life experiences of the research team are of paramount importance in this process. Due to the historical abundance of *Cycas micronesica* throughout Guam, both authors interacted with this tree throughout childhood. Additionally, after becoming

DOI: <http://doi.org/10.11609/jott.3768.9.9.10738>

Date of publication: 26 September 2017 (online & print)

Manuscript details: Ms # 3768 | Received 30 August 2017

Citation: Ramanan, S.S. (2017). Response to article on Adventitious rooting of mature *Cycas micronesica*. *Journal of Threatened Taxa* 9(9): 10738; <http://doi.org/10.11609/jott.3768.9.9.10738>

Copyright: © Ramanan 2017. Creative Commons Attribution 4.0 International License. JoTT allows unrestricted use of this article in any medium, reproduction and distribution by providing adequate credit to the authors and the source of publication.

DOI: <http://doi.org/10.11609/jott.3812.9.9.10738-10740>

Date of publication: 26 September 2017 (online & print)

Manuscript details: Ms # 3812 | Received 16 September 2017

Citation: Marler, T.E. & G.N. Cruz (2017). Best protocols for cycad propagation require more research. *Journal of Threatened Taxa* 9(9): 10738–10740; <http://doi.org/10.11609/jott.3812.9.9.10738-10740>

Copyright: © Marler & Cruz 2017. Creative Commons Attribution 4.0 International License. JoTT allows unrestricted use of this article in any medium, reproduction and distribution by providing adequate credit to the authors and the source of publication.

professionally involved in the plant sciences, we accrued a combined 32 years of studying the behavior of this and other *Cycas* species in natural and cultivated settings. A brief explanation for why we employed some methods instead of others in our recent publication is that we believed them to be the most important in fulfilling the objective. More specific reasons are detailed below.

Mist Chambers. Small adventitious stems called “pups” in the cycad nursery industry are routinely removed and nurtured to produce adventitious roots. The protocols employed in our unprecedented large-scale attempt to propagate from large apical cycad stem cuttings were primarily based on those methods, which can be accessed by any reader from multiple online resources. Based on all available knowledge, the use of intermittent mist systems would kill every experimental unit, and adding mist as a treatment in asexual propagation of cycads would not be considered by any competent research team.

Climate conditions: Cycad research is best conducted in settings that match in situ conditions. With the exception of the subpopulation on the peak of Guam’s highest elevation, all of the island’s cycad subpopulations occur in similar conditions. Our nursery was located ca. 600m west of an in situ subpopulation of *C. micronesica*. General climatic conditions of Guam can be accessed by any reader using online resources. For the duration of the propagation period, mean high temperature was 29°C, mean low temperature was 25°C, mean daily precipitation was 5.3mm, and mean daylength was 12:45hr.

Auxin and multiple nursery choices: The purpose of this adaptive management study was to determine if the protocols for asexual propagation using large stem cuttings could be scaled up to rescue the the gene pool from thousands of *C. micronesica* trees that reside within forests that will be cleared during construction activities in the near future. Using two nursery locations in the study was mandated in the project’s statement of work from the funding agency, therefore we had no choice but to include this as part of the study. Comparing dose levels of laboratory grade auxin was not considered because future Guam contractors would have difficulty obtaining the product. A retail auxin product that can be purchased locally was used to adhere to protocols that are commonly used for rooting cycad pups in the international horticulture industry. The number of experimental units needed to provide adequate replication of a dose-response treatment was not available.

Root formation on cuttings: Large cycad stem cuttings do not produce zygotic roots. A zygotic root system may



Image 1. (Left) Seedling root system developing on a *Cycas micronesica* seedlings showing a dominant taproot and subordinate lateral roots. (Right) Adventitious root system on *Cycas revoluta* stem cutting showing dominant lateral roots and no taproot.



Image 2. Adventitious root formation followed by resumption of new stem growth (blue arrow) occurred 14 months after preparing this *Cycas micronesica* pup for asexual propagation.



Image 3. Snapped *Cycas micronesica* stem sections during Guam’s recurring tropical cyclones naturally develop adventitious roots then resume stem growth (yellow arrow). The decapitated standing stem also resumes growth by developing adventitious lateral buds (blue arrows). The photograph was taken 15 months after the tropical cyclone that broke the stem.

be retained by excavating a portion of a tree's existing root system, and subsequent root growth would egress from existing root tissue that can be traced back to the original radicle that emerged from the embryo. The phenotype of a developing zygotic root system is shown in Image 1A. When large rootless stems are tended in a nursery to induce new roots, these are adventitious roots because they develop through *de novo* organogenesis from stem tissue and cannot be ontogenetically traced back to any historical root tissue. We did not bare-root any of the large rooted transplants to observe the adventitious roots, as this would have imposed unnecessary stress on the valuable transplants. However, the phenotype of an adventitious root system that developed on a *Cycas revoluta* stem cutting is shown in Image 1B. Any reader can calculate that 41% of 34 experimental units was 14 rooted transplants.

Other cycad species: No cycad horticulturist would purposefully cut off large apical stem cuttings from 49 healthy trees just to try to clone each tree. Therefore, no citations exist for other cycad species describing attempts to develop adventitious roots on large stem cuttings. This was an unprecedented opportunity to determine the success of rooting large stem cuttings from cycad trees that were located within a forested site that was destined to be destroyed for construction activities.

Additional remarks: Cycad plants are highly skilled at what they do, but they are never in a rush. When experienced horticulturists begin to grow cycad plants using methods they have developed from successfully managing common angiosperms, they often kill several specimens before learning that cycad plants often just need to be left alone. Overwatering or use of container medium that lacks adequate aeration are often the causes of these initial failures. Indeed, patience and resisting the urge to do something may be the two most important horticultural skills for managing sexual and asexual cycad propagation. Most cycad horticulturists share our experiences of patiently waiting more than one year for a cycad pup to develop adventitious roots then resume stem growth (Image 2).

No cycad horticulturist will be surprised that we were successful even though the use of large stem cuttings has never been employed for large-scale propagation. We used methods that adhered to the most important facets of asexual propagation, which included covering the exposed wound tissue with a sealant of some type, ensuring extremely high aeration within the rooting medium, respecting the need for hygiene when handling the cutting and container medium, and under-watering throughout the propagation phase. The fact that more than half of the cuttings were unsuccessful was due to the extremely poor health status of the source trees. Storage organs on healthy cycad plants contain copious levels of non-structural carbohydrates, and we believe that chronic damage by the non-native *Aulacaspis yasumatsui* armored scale infestations cause mortality by way of depleting these storage resources over time. Healthy trees with no history of herbivory damage would likely yield large stem cuttings that exhibit 100% success in developing adventitious roots. We do not envision an opportunity to verify this directly because sacrificing large healthy arborescent cycad trees just to determine if adventitious roots could be induced on large stem cuttings would be difficult to justify. But we propose that this can be studied indirectly by quantifying the response of *Cycas* stem and root non-structural carbohydrate pools to terminal infestations of *Aulacaspis yasumatsui*.

The recurrent tropical cyclones that characterize life on Guam may dislodge large structural stem sections from in situ *C. micronesica* trees. In the absence of subsequent non-native insect and mammal herbivore damage, these stems develop adventitious roots where they are in contact with the soil surface, then resume stem growth (Image 3). This natural form of cloning is evidence that cycad stems do not need anthropogenic treatments like synthetic auxins or mist chambers in order to successfully produce adventitious roots. Expanded cycad propagation research is not necessarily needed to increase cloning success rates, but is needed to speed up the process such that more established transplants can be produced in a shorter time-frame.





OPEN ACCESS



The Journal of Threatened Taxa is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at www.threatenedtaxa.org. All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use of articles in any medium, reproduction, and distribution by providing adequate credit to the authors and the source of publication.

ISSN 0974-7907 (Online); ISSN 0974-7893 (Print)

September 2017 | Vol. 9 | No. 9 | Pages: 10633–10740

Date of Publication: 26 September 2017 (Online & Print)

DOI: 10.11609/jott.2017.9.9.10633-10740

www.threatenedtaxa.org

Articles

The ecology of Harwood's Francolin *Pternistis harwoodi* (Aves: Galliformes: Phasianidae) at Merhabete District, central highlands of Ethiopia: implications for conservation
-- Abadi Mehari Abrha & Habtu Kiros Nigus, Pp. 10633–10641

Effects of the invasive Tilapia on the Common Spiny Loach (Cypriniformes: Cobitidae: *Lepidocephalichthys thermalis*) - implications for conservation
-- Sandip D. Tapkir, Sanjay S. Kharat, Pradeep Kumkar & Sachin M. Gosavi, Pp. 10642–10648

Communications

Distribution and habitat use of the endangered Dhole *Cuon alpinus* (Pallas, 1811) (Mammalia: Canidae) in Jigme Dorji National Park, western Bhutan
-- Chhimi Namgyal & Phuntsho Thinley, Pp. 10649–10655

Winter food habits of the Golden Jackal *Canis aureus* (Mammalia: Carnivora: Canidae) in Patna Bird Sanctuary, Uttar Pradesh, India
-- Khursid A. Khan, Jamal A. Khan & Narendra Mohan, Pp. 10656–10661

On the poorly-known White-spotted Skink *Lygosoma albopunctatum* (Gray, 1846) (Reptilia: Scincidae) with further topotypical records and notes on the type locality
-- S.R. Ganesh, Pp. 10662–10668

A study on the density, population structure and regeneration of Red Sanders *Pterocarpus santalinus* (Fabales: Fabaceae) in a protected natural habitat - Sri Lankamalleswara Wildlife Sanctuary, Andhra Pradesh, India
-- Chenchu Ankalaiah, Thondaladinne Mastan & Mullangi Sridhar Reddy, Pp. 10669–10674

Short Communications

First record and genetic affiliation of the Balkan Snow Vole *Dinaromys bogdanovi* (Rodentia: Cricetidae) in Albania
-- Ivan Stolarik & Daniel Jablonski, Pp. 10675–10678

Avifaunal diversity in the scrub forest of Sri Lankamalleswara Wildlife Sanctuary, Andhra Pradesh, India
-- Sumant Mali, Chelmala Srinivasulu & Asad R. Rahmani, Pp. 10679–10691

A second record of the Eastern Spadefoot Toad (Amphibia: Anura: Megophryidae: *Leptobrachium bompui* Sondhi & Ohler, 2011) with a note on its morphological variations and natural history
-- Bhaskar Saikia, Bikramjit Sinha & Ilona J. Kharkongor, Pp. 10692–10696

Spiders of Gujarat: a preliminary checklist
-- Archana Yadav, Reshma Solanki, Manju Siliwal & Dolly Kumar, Pp. 10697–10716

An ethnomycological survey of Jaunsar, Chakrata, Dehradun, India
-- Manoj Kumar, N.S.K. Harsh, Rajendra Prasad & Vijay Vardhan Pandey, Pp. 10717–10725

Notes

First record of Rufous-tailed Scrub Robin *Cercotrichas galactotes* (Aves: Passeriformes: Muscicapidae) from Jammu & Kashmir, India
-- Neeraj Sharma, Pp. 10726–10728

Sightings of the Great White Pelican *Pelecanus onocrotalus* (Linn. 1758) (Aves: Pelicaniformes: Pelicanidae) in Pallikarai Marshlands, Tamil Nadu, India
-- Kannan Thirunaranan, Samidurai Jayakumar, Sankaranathan Sivaraman & Santhanakrishnan Babu, Pp. 10729–10732

A note on three interesting Heteroptera from Kolhapur, Maharashtra, India
-- S.M. Gaikwad, Y.J. Koli & G.P. Bhawane, Pp. 10733–10737

Response & Reply

Response to article on Adventitious rooting of mature *Cycas micronesica*
-- S. Suresh Ramanan, P. 10738

Best protocols for cycad propagation require more research
-- Thomas Edward Marler & Gil Naputi Cruz, Pp. 10738–10740