

OPEN ACCESS

The Journal of Threatened Taxa is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at www.threatenedtaxa.org. All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use of articles in any medium, reproduction, and distribution by providing adequate credit to the authors and the source of publication.



Journal of Threatened Taxa

Building evidence for conservation globally

www.threatenedtaxa.org

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

NOTE

FORAGING HABITS OF THE RED FOX *VULPES VULPES* (MAMMALIA: CARNIVORA: CANIDAE) IN THE HIMALAYA, INDIA

Aishwarya Maheshwari

26 September 2018 | Vol. 10 | No. 10 | Pages: 12418–12421
10.11609/jott.3968.10.10.12418-12421



For Focus, Scope, Aims, Policies and Guidelines visit <http://threatenedtaxa.org/index.php/JoTT/about/editorialPolicies#custom-0>
For Article Submission Guidelines visit <http://threatenedtaxa.org/index.php/JoTT/about/submissions#onlineSubmissions>
For Policies against Scientific Misconduct visit <http://threatenedtaxa.org/index.php/JoTT/about/editorialPolicies#custom-2>
For reprints contact info@threatenedtaxa.org

Partners



صندوق محمد بن زايد
للمحافظة على
الكائنات الحية
The Mohamed bin Zayed
SPECIES CONSERVATION FUND



zoo!
ZÜRICH

Member



Publisher & Host



Threatened Taxa



ISSN 0974-7907 (Online)
ISSN 0974-7893 (Print)

OPEN ACCESS



**FORAGING HABITS OF THE RED FOX
VULPES VULPES (MAMMALIA: CARNIVORA:
CANIDAE) IN THE HIMALAYA, INDIA**

Aishwarya Maheshwari 

Banda University of Agriculture and Technology, Near New Circuit House, Banda, Uttar Pradesh 210001, India
aishwaryamaheshwari@gmail.com

The Red Fox *Vulpes vulpes* is one of the most widely distributed and extremely adaptable carnivore in the world (Macdonald & Reynolds 2004), and found in a variety of habitats ranging from the arctic to temperate deserts (Macdonald & Reynolds 2004). It is an omnivorous opportunistic predator and feeds on essentially anything easily available or small enough to catch, from the wilderness to cities (Harris & Smith 1987; Jędrzejewski & Jędrzejewska 1992; Scott et al. 2014). Small mammals, birds and insects are the major food materials recorded in the Red Fox diet (Goszczynski 1974; Meisner et al. 2014). Red Foxes are mostly nocturnal (Ables 1969; Macdonald 1980; Travainiet et al. 1993; Weber et al. 1994) but their activity pattern and movement may overlap with the availability of forage and level of disturbance (Macdonald 1980; Lovari et al. 1994; König 2008). Keeping this in view, the present study is an attempt to understand if the nocturnal behavior of the Red Fox alters due to the easy availability of food resources in the daytime.

The study was conducted in 12 villages covering approximately 1,000km² of Kargil District with an area of about 14,000km² (Fig. 1). Kargil is a mountainous cold desert in Ladakh region with little or sparse vegetation and represents the biogeographic zone 1B (Trans-Himalaya-Tibetan Plateau) of India (Rodgers et al. 2000). The general elevation of Kargil ranges from 2,934–7,410 m with an average elevation of 3,400m (Maheshwari 2016).

Observations on the Red Fox were recorded during field studies on snow leopards *Panthera uncia* and associated species with special reference to large carnivore-human conflict, conducted from April 2009 to November 2012 (Maheshwari 2016). Due to the topography and remoteness of the area, all fieldwork was carried out in the form of discrete field expeditions that involved camping in the different areas. Each field survey usually lasted 10–15 days. Altogether, 1,100km were traversed on foot covering an altitudinal zone of 3,000–5,200 m. Every sighting of the Red Fox was recorded during the fieldwork, and interviews were conducted of all the 664 households across 12 villages in the study landscape. Information was gathered on livestock predation such as species and number of attacks with time and place of attack by Red Fox during the study period. While collecting data on Red Fox predation on livestock in order to reduce probability of response bias and avoid overestimation of livestock predation, protocols under participatory rural appraisal (PRA) (Maheshwari et al. 2014) were employed by using the semi-structured interview technique of PRA.

DOI: <https://doi.org/10.11609/jott.3968.10.10.12418-12421>

Editor: L.A.K. Singh, Bhubaneswar, Odisha, India.

Date of publication: 26 September 2018 (online & print)

Manuscript details: Ms # 3968 | Received 22 December 2017 | Final received 05 June 2018 | Finally accepted 20 August 2018

Citation: Maheshwari, A. (2018). Foraging habits of the Red Fox *Vulpes vulpes* (Mammalia: Carnivora: Canidae) in the Himalaya, India. *Journal of Threatened Taxa* 10(10): 12418–12421; <https://doi.org/10.11609/jott.3968.10.10.12418-12421>

Copyright: © Maheshwari 2018. Creative Commons Attribution 4.0 International License. JoTT allows unrestricted use of this article in any medium, reproduction and distribution by providing adequate credit to the authors and the source of publication.

Funding: The Rufford Foundation.

Competing interests: The author declares no competing interests.

Acknowledgements: I am grateful to Rufford Foundation for financially supporting the work. The Department of Wildlife Protection, Jammu & Kashmir is particularly thanked for their permission, interest and support during the work. I thank my colleagues at the Wildlife Institute of India, Dehradun and Banda University of Agriculture and Technology, Banda for their continuous encouragement and support while writing the manuscript. I am thankful to Kazim for his continuous assistance during the fieldwork. I extend my gratitude to Mukesh and anonymous reviewers for their valuable feedback in improving the manuscript.






Figure 1. Study Area: location of the 12 sampled villages in Kargil, Ladakh, India

With the help of field assistants data on visual encounters ($n=220$) of the Red Fox from April 2009 to November 2012 were gathered. The maximum sightings were recorded during 15.01 to 18.00hr (45.4%) followed by 12.01 to 15.00hr (25.9%), 09.01 to 12.00hr (25.4%) and 06.00 to 09.00hr(13.2%). To understand Red Fox movement during day-light hours in a human dominated landscape, data were also collected on the availability of food resources, e.g., free ranging domestic fowls and inattentive young ones of the sheep/ goats in the villages (Image 1). A total of 230 domestic fowls and 74 young ones of the sheep/ goats were reportedly killed by the Red Fox in Kargil. Of 12 villages, the highest livestock attacks were recorded in Sapi (15.8%) followed by the remaining 11 villages (Table 1).

Data obtained from locals on the time of predation was overlapping with the time of sighting of the Red Fox in the villages. Most (40.9%) of the domestic fowls and 35.1% of sheep and goats' predation events were recorded during 09:01–12:00 hr followed by 34.3% (domestic fowls) and 27% (sheep and goats) during 12:01–15:00 hr (Table 2).

During the day time, when most of the family members were engaged in domestic work and other livestock grazing, they set free the fowls and young ones of the sheep and goats to move on their own and feed upon freely and naturally accessible food. At this point they were vulnerable for predation by the Red Fox.

Foxes may be found during the day pursuing prey and resting (Meisner et al. 2014). One breed of fox that is definitely diurnal is the Island Fox *Urocyon littoralis* (U.S. Fish & Wildlife Service 2015). Red Foxes are typically nocturnal animals, especially inhabiting in and around urban areas, to avoid being seen or disturbed by humans (Scott et al. 2014). Local people report that the Red Fox has got accustomed to raiding villages and houses for food. As it takes a lot of energy to scare these foxes, some locals are retaliating by killing the Red Foxes. Anthropogenic feeding has been reportedly supporting an increase in density of Red Fox range from 2–30 adults/km² (Baker et al. 2000; Soulsbury et al. 2010; Scott et al. 2014) in the urban areas. But in Kargil, where local communities are primarily agro-pastoral and livestock rearing is one of the major sources of livelihood, loss of

Table 1. Red Fox predation on domestic fowls and sheep/ goats across 12 villages in Kargil

	Name of village	Total number of fowl predation	Total number of sheep and goat predation
1	Bartoo	14	4
2	Pangbar	7	3
3	Yarkashing	15	4
4	Bilching	4	2
5	Umba	30	9
6	Ichoo	12	3
7	Mulbek	22	7
8	Shergandi	12	1
9	Fokar	26	9
10	Kanji	28	12
11	Sapi	38	10
12	Wakhah	22	10
Total		230	74

Table 2. Red Fox predation during different time intervals of the day

Time interval (hr)	Domestic fowls	Sheep and goats	Total
06:00–09:00	30 (13.0%)	14 (18.9%)	44 (14.5%)
09:01–12:00	94 (40.9%)	26 (35.1%)	120 (39.5%)
12:01–15:00	79 (34.3%)	20 (27.0%)	99 (32.6%)
15:01–18:00	27 (11.7%)	14 (18.9%)	41 (13.5%)
Total numbers	230	74	304

livestock leads to retaliatory killing (Maheshwari 2016). Thus, similar to other carnivores in the global scenario, the Red Fox is also a victim of retaliation in Kargil. During the study five cases of retaliation against the Red Fox were recorded. Besides, there were two records of road kills during the same period (Image 2). In the absence of a proper mechanism to monitor and record Red Fox killing at landscape level this study presents only a fraction of the actual cases of retaliation and road kills. Nevertheless, retaliation and road kills constitute the major threats to the overall survival of the Red Fox population.

The lack of livestock guarding practices and poor or no search efforts by people to locate 'missing' animals are two of the major factors responsible for livestock loss in Kargil. The loss of domestic fowls and livestock constitute one of the major threats to the rural economy and the Red Fox is one of the major predators in Kargil. Diurnal alteration in foraging behavior of the Red Fox could be due to competition with the Snow Leopard

**Image 1. Red Fox predation on domestic fowl****Image 2. Red Fox road kill**

and Wolf *Canis lupus chanco* (Maheshwari 2016). About 8.3% livestock loss (2009–2012) was due to predation by large carnivores, i.e., a total of 1113 heads of livestock were reportedly killed by wolf (43.6%) followed by unknown predators (31.4%) and Snow Leopard (21.5%) in the study site, which comes to 2.8% of total annual livestock losses (Maheshwari 2016). This study adds to the limited information available on the dynamics of human-Red Fox interaction in Kargil.

The Red Fox is a well-studied species across the world (Macdonald & Reynolds 2004), but information regarding its distribution, ecology, and subspecies remains rather limited in India (Maheshwari et al. 2013). The species in India, which is relatively better studied among the fox species, is the Indian Fox *Vulpes bengalensis* (Home 2005; Kumara & Singh 2012; Maurya et al. 2012). One of the important notes for future studies could be to gather crucial baseline information on the status, distribution, ecology and interaction with human of the other fox species and subspecies (Maheshwari et al. 2013) in India. This is crucial in quantifying changes in Red Fox densities due to interface with anthropogenic dimensions, and

develop strategies for conservation management.

References

- Ables, E.D. (1969).** Activity studies of Red Foxes in southwestern Wisconsin. *Journal of Wildlife Management* 33: 145-153.
- Baker, P.J., S.M. Funk, S. Harris & P.C. White (2000).** Flexible spatial organization of urban foxes, *Vulpes vulpes*, before and during an outbreak of sarcoptic mange. *Animal Behavior* 59: 127-146.
- Goszczynski, J. (1974).** Studies on the food of foxes. *Acta Theriologica* 19: 1-18.
- Harris, S. & G.C. Smith (1987).** The use of sociological data to explain the distribution and numbers of Urban Foxes (*Vulpes vulpes*) in England and Wales. *Symposium of Zoological Society London* 58: 313-328.
- Home, C. (2005).** Resource Utilization by Indian Fox (*Vulpes bengalensis*) in Kutch, Gujarat. MSc dissertation, Wildlife Institute of India, Dehradun, India.
- Jędrzejewski, W. & B. Jędrzejewska (1992).** Foraging and diet of the Red Fox *Vulpes vulpes* in relation to variable food resources in Białowieża National Park, Poland. *Ecography* 15: 212-220.
- König, A. (2008).** Fears, attitudes and opinions of suburban residents with regards to their urban foxes. *European Journal of Wildlife Research* 54: 101-109.
- Kumara, H.N. & M. Singh (2012).** Distribution, den characteristics and diet of the Indian Fox *Vulpes bengalensis* (Mammalia: Canidae) in Karnataka, India: preliminary observations. *Journal of Threatened Taxa* 4(14): 3349-3354; <https://doi.org/10.11609/JoTT.o3046.3349-54>
- Lovari, S., P. Valier & M.R. Lucchi (1994).** Ranging behavior and activity of Red Foxes (*Vulpes vulpes*: Mammalia) in relation to environmental variables, in a Mediterranean mixed pinewood. *Journal of Zoology* 232: 323-339.
- Macdonald, D.W. (1980).** The Red Fox, *Vulpes vulpes*, as a predator upon earthworms, *Lumbricus terrestris*. *Zeitschrift für Tierpsychologie* 52: 171-200.
- Macdonald, D.W. & J.C. Reynolds (2004).** Red Fox *Vulpes vulpes* Linnaeus, 1758. pp. 129-134. In: Sillero-Zubiri, C., M. Hoffmann & D.W. Macdonald (eds.) *Canids: Foxes, Wolves, Jackals and Dogs. Status Survey and Conservation Action Plan*. IUCN/SSC Canid Specialist Group. Gland, Switzerland and Cambridge, UK.
- Maheshwari, A. (2016).** Conservation and management of Snow Leopard and co-predators with special reference of large carnivore-human conflicts in the select areas of western Himalayas. PhD Thesis, Saurashtra University, Rajkot, Gujarat, India.
- Maheshwari, A., N. Midha, A. Paliwal, B. Sharma, P.S. Ghose, P. Shreshtha & S. Paranjpe (2013).** Note on the subspecies of Red Fox and Tibetan Sand Fox: Preliminary overview from the Himalayas, India. *Journal of the Bombay Natural History Society* 110: 193-196.
- Maheshwari, A., N. Midha & A. Cherukupalli (2014).** Participatory rural appraisal and compensation intervention: challenges and protocols while managing large carnivore-human conflict. *Human Dimensions of Wildlife* 19: 62-71.
- Maurya, K.K., I.P. Bopanna & Y.V. Jhala (2012).** Estimating food intake from scats in the omnivorous Indian Fox (*Vulpes bengalensis*). *Journal of the Bombay Natural History Society* 109: 177-181.
- Meisner, K., P. Sunde, K.K. Clausen, P. Clausen, C.C., Fælled & M. Hoelgaard (2014).** Foraging ecology and spatial behaviour of the Red Fox (*Vulpes vulpes*) in a wet grassland ecosystem. *Acta Theriologica* 59: 377-389.
- Rodgers, W.A., H.S. Panwar & V.B. Mathur (2000).** Wildlife Protected Area Network in India: A Review (Executive Summary). Wildlife Institute of India, Dehradun.
- Scott, D.M., M.J. Berg, B.A. Tolhurst, A.L.M. Chauvenet, G.C. Smith, K. Neaves, J. Lochhead & P.J. Baker (2014).** Changes in the distribution of Red Foxes (*Vulpes vulpes*) in urban areas in Great Britain: findings and limitations of a media-driven nationwide survey. *PLoS ONE* 9: e99059; <https://doi.org/10.1371/journal.pone.0099059>
- Soulsbury, C.D., P.J. Baker, G. Iossa & S. Harris (2010).** Red Foxes (*Vulpes vulpes*), pp 63-75. In: Gehrt, S.D., S.P.D. Riley & B.L. Cypher (eds.). *Urban Carnivores: Ecology, Conflict, and Conservation*. Johns Hopkins University Press, Baltimore, Md.
- Travaini, A., J.J. Aldama, R. Laffitte & M. Delibes (1993).** Home range and activity patterns of Red Fox *Vulpes vulpes* breeding females. *Acta Theriologica* 38: 427-434.
- U.S. Fish & Wildlife Service (2015).** Recovery plan for four subspecies of Island Fox (*Urocyon littoralis*). U.S. Fish and Wildlife Service, Sacramento, California.
- Weber, J.M., S. Meia & S. Aubry (1994).** Activity of foxes, *Vulpes vulpes* in the Swiss Jura mountains. *Zeitschrift für Säugetierkunde* 59: 9-13.





OPEN ACCESS



The Journal of Threatened Taxa is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at www.threatenedtaxa.org. All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use of articles in any medium, reproduction, and distribution by providing adequate credit to the authors and the source of publication.

ISSN 0974-7907 (Online); ISSN 0974-7893 (Print)

September 2018 | Vol. 10 | No. 10 | Pages: 12299–12442

Date of Publication: 26 September 2018 (Online & Print)

DOI: 10.11609/jott.2018.10.10.12299-12442

www.threatenedtaxa.org

Communications

Urban biodiversity: an insight into the terrestrial vertebrate diversity of Guwahati, India

-- Jayaditya Purkayastha, Pp. 12299–12316

Status of raptors in the Moyar River Valley, Western Ghats, India

-- N.R. Anoop, S. Babu, S. Bharathidasan & R. Nagarajan, Pp. 12317–12327

Species composition and abundance estimates of reptiles in selected agroecosystems in southern Western Ghats, India

-- Abhirami Mini Jayakumar & Paingamadathil Ommer Nameer, Pp. 12328–12336

Comparison of beach profiles conducive for turtle nesting in Andaman

-- Subramanian Narayani, Sasidharan Venu & Andrea Joan D'Silva, Pp. 12337–12343

Short Communications

A new record of the rare Hardwicke's Woolly Bat *Kerivoula hardwickii* (Horsfield, 1824) (Mammalia: Chiroptera: Vespertilionidae) after 23 years from a lowland rainforest of Sri Lanka

-- Dinesh Gabadage, Gayan Edirisinghe, Madhava Botejue, Kalika Perera, Thilina Surasinghe & Suranjan Karunarathna, Pp. 12344–12349

Alarming population status of the Grizzled Giant Squirrel *Ratufa macroura* (Mammalia: Rodentia: Sciuridae) in Chinnar Wildlife Sanctuary, the Western Ghats, India

-- Kiran Thomas & Paingamadathil Ommer Nameer, Pp. 12350–12356

Distribution and population status of Sambar *Rusa unicolor* (Mammalia: Cetartiodactyla: Cervidae) from Aravalli landscape with a note on its first record from Aravalli Hills of Haryana, India

-- Paridhi Jain, Anchal Bhasin, Gautam Talukdar & Bilal Habib, Pp. 12357–12362

Delayed peracute capture myopathy in a Himalayan Ibex *Capra sibirica* (Mammalia: Cetartiodactyla: Bovidae)

-- Umar Nazir Zahid, Latief Mohammad Dar, Umar Amin, Showkat Ahmad Shah, Rashid Yahya Naqash, Dil Mohammed Makhdoomi, Shayuaib Ahmad Kamil & Intesar Suhail, Pp. 12363–12367

Checklist of the avifauna of Sagareshwar Wildlife Sanctuary, Maharashtra, India

-- Sharad Datt Apte, Vijay Bhagwan Tuljapurkar & Girish Avinash Jathar, Pp. 12368–12375

The rediscovery of Rurk's Cat Skink *Ristella rurkii* Gray, 1839 (Reptilia: Ristellidae) with remarks on distribution and natural history

-- Sumaithangi Rajagopalan Ganesh, Pp. 12376–12381

Dietary assessment of five species of anuran tadpoles from northern Odisha, India

-- Syed Asrafuzzaman, Susmita Mahapatra, Jasmin Rout & Gunanidhi Sahoo, Pp. 12382–12388

Partners



Inventory of prong-gilled mayflies (Ephemeroptera: Leptophlebiidae) of India with records of endemic taxa

-- C. Selvakumar, Kailash Chandra & K.G. Sivaramakrishnan, Pp. 12389–12406

First record of a coreid bug *Anhomoeus fusiformis* Hsiao (Hemiptera: Heteroptera: Coreidae: Coreinae: Anhomoeini) from India

-- Sadashiv V. More & Hemant V. Ghate, Pp. 12407–12412

The gilled mushroom *Amanita spissacea* (Amanitaceae): a new report for India

-- Hmar Lalrinawmi, John Zothanzama, Benjamin W. Held, Josiah M.C. Vabeikokhei, Zohmangaiha & Robert A. Blanchette, Pp. 12413–12417

Notes

Foraging habits of the Red Fox *Vulpes vulpes* (Mammalia: Carnivora: Canidae) in the Himalaya, India

-- Aishwarya Maheshwari, Pp. 12418–12421

First record of Yellow-Rumped Flycatcher *Ficedula zanthopygia* (Hay, 1845) (Aves: Passeriformes: Muscipidae) in eastern India

-- Manaranjan Das & Subrat Debata, Pp. 12422–12424

Additional field records provide further resolution of the distribution of the Water Monitor *Varanus salvator* (Squamata: Varanidae) in northwestern Myanmar

-- Steven G. Platt, Myo Min Win & Thomas R. Rainwater, Pp. 12425–12428

The first record of The Blue Admiral *Kaniska canace* Linnaeus, 1763 (Nymphalidae: Lepidoptera) from Bangladesh

-- Amit Kumer Neogi, Md Jayedul Islam, Md Shalauddin, Anik Chandra Mondal & Safayat Hossain, Pp. 12429–12431

First record of *Hislopia malayensis* Annandale, 1916 (Bryozoa: Gymnolaemata) from freshwaters of India

-- Ananta Dnyanoba Harkal & Satish Sumanrao Mokashe, Pp. 12432–12433

An extended distribution record of Western Ghats species *Litsea oleoides* (Meissn.) Hook.f. (Lauraceae) from Matheran, Maharashtra, India

-- Radha Veach & Gurumurthi Hegde, Pp. 12434–12438

Notes on *Jasminum andamanicum* N.P. Balakr. & N.G. Nair (Oleaceae) from Andaman & Nicobar Islands, India

-- P. Murugan & K. Karthigeyan, Pp. 12439–12441

Miscellaneous

National Biodiversity Authority

Member



Publisher & Host

