

The Journal of Threatened Taxa (JoTT) is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at www.threatenedtaxa.org. All articles published in JoTT are registered under Creative Commons Attribution 4.0 International License unless otherwise mentioned. JoTT allows allows unrestricted use, reproduction, and distribution of articles in any medium by providing adequate credit to the author(s) and the source of publication.

Journal of Threatened Taxa

Building evidence for conservation globally

www.threatenedtaxa.org ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

NOTE POLLINATORS OF SIKKIM MANDARIN ORANGE *CITRUS RETICULATA* (SAPINDALES: RUTACEAE)

Urbashi Pradhan & M. Soubadra Devy

26 March 2019 | Vol. 11 | No. 5 | Pages: 13625–13628 DOI: 10.11609/jott.4528.11.5.13625-13628





For Focus, Scope, Aims, Policies, and Guidelines visit https://threatenedtaxa.org/index.php/JoTT/about/editorialPolicies#custom-0 For Article Submission Guidelines, visit https://threatenedtaxa.org/index.php/JoTT/about/submissions#onlineSubmissions For Policies against Scientific Misconduct, visit https://threatenedtaxa.org/index.php/JoTT/about/editorialPolicies#custom-2 For reprints, contact <ravi@threatenedtaxa.org>

The opinions expressed by the authors do not reflect the views of the Journal of Threatened Taxa, Wildlife Information Liaison Development Society, Zoo Outreach Organization, or any of the partners. The journal, the publisher, the host, and the partners are not responsible for the accuracy of the political boundaries shown in the maps by the authors.

Partner A conservation Partner Conservation Partner Partn

Member







POLLINATORS OF SIKKIM MANDARIN ORANGE *CITRUS RETICULATA* (SAPINDALES: RUTACEAE)

Urbashi Pradhan¹ & M. Soubadra Devy²

^{1.2} Ashoka Trust for Research in Ecology and the Environment (ATREE), Royal Enclave, Srirampura, Jakkur, Bengaluru, Karnataka 560064, India.
¹ Manipal Academy of Higher Education, Tiger Circle Road, Madhav

Nagar, Manipal, Karnataka 576104, India

¹urbashipradhan@gmail.com (corresponding author),

² soubadra@atree.org

Sikkim Mandarin Orange (*Citrus reticulata* Blanco, 1837) is a member of the Rutaceae family and a commercially desirable variety of the mandarin group native to Sikkim. The Sikkim Mandarin Orange (SMO) growing area lies at an altitudinal range of 700–1,500 m and it is an annual flowering plant. Mandarin orange is dependent on bees for its pollination and pollinators help in higher yield and increased fruit set (ICIMOD 2003). Irrespective of large cardamom yield decline due to pollinator deficiency in Sikkim (Sinu & Shivanna 2007), till date there exists no systematic study on the range of pollinators for SMO. This study aims to bridge this gap especially when a large proportion of farmers are dependent on the SMO for cash income.

Our study area spanned the East, West and South districts of Sikkim. The southern part of the state, which lies in the altitude range of 600–1500 m provides an ideal climate for SMO cultivation (DHCCD 2015). Data was collected across 72 SMO orchards from 2011

to 2013. These orange orchards were selected within an altitudinal gradient of 700–1,452 m and were spread across 316km² (Fig. 1).

Pollinator visitation: At each site, 150 flowers were tagged and observed from 08.00–17.00 h to record insect species that visit them. Intra-floral foraging behavior of each insect species was carefully



ISSN 0974-7907 (Online) ISSN 0974-7893 (Print)



observed to note whether it is a pollinator or a forager. SMO bear self-fertile, bisexual flowers and pollen movement is facilitated by pollinators. Transparent plastic bags were used to trap insects visiting the flowers to avoid any fruit loss during their collection. Collected samples were preserved in 70% ethanol and subsequently identified in the laboratory. Insects which were not seen touching the flower reproductive parts were not collected for identification.

We recorded 24 species of insects during the study period (2011–2013). Common Honey Bee *Apis cerana* was the most dominant pollinator followed by hoverflies belonging to eight genera, namely, *Episyrphus* sp., *Melanostoma* sp., *Ischiodon* sp., *Eristalis* sp., *Eristalinus* sp., *Scaeva* sp., *Episyrphus* sp., and *Eupeodes* sp. (Image 1,2). This was followed by stingless bees (Hymenoptera), seed bug (Hemiptera), and beetles (Coleoptera) that were sparse visitors. Recorded insects were both pollen and nectar feeders. Bees (Hymenoptera) and hoverflies (Diptera) visited flowers in groups while most of the

DOI: https://doi.org/10.11609/jott.4528.11.5.13625-13628 | ZooBank: urn:lsid:zoobank.org:pub:8120DF28-84C5-48D0-B1DF-2F6C8F8312C3

Editor: Kannan C.S. Warrier, Institute of Forest Genetics and Tree Breeding, Coimbatore, India.

Manuscript details: #4528 | Received 28 August 2018 | Final received 06 February 2019 | Finally accepted 16 March 2019

Citation: Pradhan, U. & M.S. Devy (2018). Pollinators of Sikkim Mandarin Orange Citrus reticulata (Sapindales: Rutaceae). Journal of Threatened Taxa 11(5): 13625–13628. https://doi.org/10.11609/jott.4528.11.5.13625-13628

Copyright: © Pradhan & Devy 2019. Creative Commons Attribution 4.0 International License. JoTT allows unrestricted use, reproduction, and distribution of this article in any medium by adequate credit to the author(s) and the source of publication.

Funding: Department of Science and Technology, Government of India; National Mission for Himalayan Studies (NMHS) under the Ministry of Environment, Forest and Climate Change, Government of India (NMHS/2015-16/HF11/11).





Date of publication: 26 March 2019 (online & print)

Competing interests: The authors declare no competing interests.

Acknowledgements: We would like to thank Dr Belavadi, Dr Yashwanth and Dr Arathi at GKVK, Bangalore and Dr Kumar Ghorpade, at University of Agricultural Science, Dharwad for helping us identify our pollinator samples. We are immensely grateful to the people of the study villages for their hospitality and support. The officials of the Department of Horticulture and Cash Crop, especially Mr K.K. Singh, ex- Principal Secretary, Mr Khorlo Bhutia, the present Principal Secretary, Mr P.T. Bhutia and field staffs were all generously helpful. We would like to thank the Forest Department officials, PCCF Mr S.T.Lachungpa and PCCF Dr Thomas Chandy, Mrs Usha Lachungpa (Ex- Scientific officer) for giving the necessary permission to work in Sikkim. We take this opportunity to thank the donors—Department of Biotechnology, Government of India and National Mission on Himalayan Studies(NMHS), The Ministry of Environment, Forest & Climate Change (MoEF&CC)—who generously funded this research.



Figure 1. Location of orchards (in red dots) within the northeastern Indian state of Sikkim.

beetles and seed bugs visited individually. All the insects landed on the petal and foraged for pollen placed on top of the flower and nectar at the flower base. In this process all the insects invariably touch both anther and stigma of SMO flower. An insect visitor was called a pollinator when the ventral side of insect's body containing pollen load touched the reproductive part of flowers.

SMO is an evergreen plant showing flowering response to early monsoon shower starting in the mid of February. Flowering period lasts for a month, from late February to early April. Orchards in the lower altitude starts flowering earlier followed by orchards in the higher altitudes. Flowers are white in colour with strong scent attracting a range of insects for pollination. Highlighting the importance of pollinators of the mandarin orange, in a study conducted by the International Center for Integrated Mountain Development (2003), pollination was seen to increase the yield of mandarin orange by four times compared to pollinator excluded flowers. Honey bee (*Apis* sp.) has been reported as a major pollinator of different varieties of *Citrus* sp. from across the world, for example, Mandarin orange *Citrus reticulata* in Nepal

is pollinated by A. cerana, A. dorsata, A. florea, and A. mellifera (International Center for Integrated Mountain Development 2003). Kinnow Citrus reticulata, a hybrid between mandarin orange and sweet lime, was reported to be pollinated by A. dorsata and A. florea in Pakistan (Manzoorul-Haq et al. 1978). Results of our study show only A. cerana visited mandarin orange flowers, while A. dorsata or A. florea, which were recorded in other studies, were not observed even outside our experiment sites during the study period. Hoverflies, although not reported as pollinators of mandarin oranges earlier, are known to pollinate rapeseed oil (Jauker & Wolters 2008), apple (Solomon & Kendall 1970), and strawberries (Kendall et al. 1971). Both bees and flies visited flowers in groups and visited more than one flower at a time, possibly aiding in cross/sexual pollination (Raju et al. 2012). Visits by other taxa such as butterflies, stingless bees, and beetles to orange flowers were less in comparison to bees and flies. However, the importance of these wild pollinators in sustaining pollination of SMO needs further exploration.

Pollinators of Sikkim Mandarin Orange

Table 1	. Pollinators of	ⁱ Sikkim Mand	larin Orange
---------	------------------	--------------------------	--------------

	Order	Family	Sub family	Genus	Species	Altitude range	Forage collected
1	Coleoptera	Coccinellidae	Coccinellidae	Oenopia	kirbyi (Mulsant)	700–1400	nectar + pollen
2		Scarabaeidae	Rutelinae	Anomala	sp.	700–1400	nectar + pollen
3		Scarabaeidae	Citoniinae	Clinteria	sp.	700–1400	nectar + pollen
4		Chrysomelidae	Eumolpinae	Chrysonopa	sp.	700–1400	nectar + pollen
5		Chrysomelidae	Galerucinae	Galerucinae	sp.	700–1400	nectar + pollen
6	Diptera	Calliphoridae	Chrysomyinae	Chrysomya	sp.	1000–1400	nectar + pollen
7		Rhiniidae	Rhiniinae	Rhinia	sp.	0800-1400	nectar + pollen
8		Sarcophagidae	Paramacronychiinae	Wohlfartia	sp.	0800-1400	nectar + pollen
9		Syrphidae	Syrphinae	Episyrphus	sp.	0800-1400	nectar + pollen
10		Syrphidae	Syrphinae	Melanostoma	sp.	900–1400	nectar + pollen
11		Syrphidae	Syrphinae	Ischiodon	scutellaris (Fabricius)	0800-1400	nectar + pollen
12		Syrphidae	Eristalinae	Eristalis	<i>tenax</i> (Linnaeus)	0800-1400	nectar + pollen
13		Syrphidae	Syrphinae	Scaeva	<i>pyrastri</i> (Linnaeus)	900–1400	nectar + pollen
14		Syrphidae	Syrphinae	Eupeodes	confrater (Wiedemann)	900–1400	nectar + pollen
15		Syrphidae	Eristalinae	Eristalinus	taeniops (Wiedemann)	900–1400	nectar + pollen
16		Syrphidae	Syrphinae	Episyrphus	Viridaureus	900–1400	nectar + pollen
17		Syrphidae	Eristalinae	Eristalis	basifemorata(Brunetti)	700–1400	nectar + pollen
18	Hemiptera	Lygaeidae	Lygaeinae	Spilostethus	pandurus (Scopoli)	700–1400	nectar + pollen
19		Lygaeidae	Lygaeinae	Graptostethus	incertus (Walker)	700–1400	nectar + pollen
20		Largidae	Physopeltinae	Physopelta	<i>gutta gutta</i> (Burmeister)	700–1400	nectar + pollen
21	Hymenoptera	Halictidae	Halictinae	Seladonia sp	sp.	700–1200	nectar + pollen
22		Halictidae	Halictinae	Lasioglossum	sp.	700–1200	nectar + pollen
23		Apidae	Apinae	Apis	cerana	700–1500	nectar + pollen
24		Apidae	Apinae	Tetragonula	sp.	800-1200	nectar + pollen



Image 1. Apis cerana pollinating Sikkim Mandarin Orange flower.



Image 2. *Apis cerana* and *Eristalis* sp. visiting Sikkim Mandarin Orange flowers.

Pollinators of Sikkim Mandarin Orange

References

- DHCCD (2015). AgriSkm. Available from: http://sikkimagri.gov.in/ General/Eng/CropsDetails.aspx?ID=23. Accessed on 22 September 2015.
- **ICIMOD** (2003). A report: Cash Crop Farming in Nepal: The importance of pollinator diversity and managed pollination in *Citrus*. International Center for Integrated Mountain Development Jawalakhel, Kathmandu, Nepal, 49pp
- Jauker, F. & V. Wolters (2008). Hover flies are efficient pollinators of oilseed rape. *Oecologia* 156: 819–823. https://doi.org/10.1007/ s00442-008-1034-x
- Kendall, D.A., D. Wilson, C.G. Guttridge & H.M. Anderson (1971). Testing Eristalis as a pollinator of covered crops. *Long Ashton Res Stn Rep* 1971: 120–121.
- Manzoorul-Haq, M. Rafie-ul-Din & A. Ghaffar (1978). Effect of insect pollination on fruit bearing in Kinnow mandarin (*Citrus reticulata*), and physical and chemical properties of the fruit. *Journal of Apicultural Research* 17: 47–49.
- Raju, A.J.S., P.V.S. Rao, R. Kumar & S.R. Mohan (2012). Pollination biology of the crypto-viviparous Avicennia species (Avicenniaceae). Journal of Threatened Taxa 4: 3377–3389. https://doi.org/10.11609/ JoTT.o2919.3377-89
- Sinu, P.A. & K.R. Shivanna (2007). Pollination biology of large cardamom (*Amomum subulatum*). Current Science 93: 548–552.
- Solomon, M.E. & D.A. Kendall (1970). Pollination by the syrphid fly, *Eristalis tenax*. Available online at http://agris.fao.org/agris-search/ search.do?recordID=XE20122001901. Accessed on 17 September 2015.



Journal of Threatened Taxa | www.threatenedtaxa.org | 26 March 2019 | 11(5): 13625–13628





The Journal of Threatened Taxa (JoTT) is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at www.threatenedtaxa.org. All articles published in JoTT are registered under Creative Commons Attribution 4.0 International License unless otherwise mentioned. JoTT allows allows unrestricted use, reproduction, and distribution of articles in any medium by providing adequate credit to the author(s) and the source of publication.

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

March 2019 | Vol. 11 | No. 5 | Pages: 13511–13630 Date of Publication: 26 March 2019 (Online & Print) DOI: 10.11609/jott.2019.11.5.13511-13630

www.threatenedtaxa.org

Article

Factors affecting diversity and distribution of threatened birds in Chitwan National Park, Nepal

– Jagan Nath Adhikari, Bishnu Prasad Bhattarai & Tej Bahadur Thapa, Pp, 13511–13522

Communications

Encounter rates and group sizes of diurnal primate species of Mole National Park, Ghana – Edward Debrah Wiafe, Pp. 13523–13530

Estimating Leopard Panthera pardus fusca (Mammalia: Carnivora:

Felidae) abundance in Kuno Wildlife Sanctuary, Madhya Pradesh, India

 Devavrat Pawar, Howard P. Nelson, Divya R.L. Pawar & Sarika Khanwilkar, Pp. 13531–13544

Food composition of Indian Eagle Owl *Bubo bengalensis* Franklin (Aves: Strigiformes: Strigidae) from Tiruchirappalli District, Tamil Nadu, India

 Tamilselvan Siva, Periyasamy Neelanarayanan & Vaidyula Vasudeva Rao, Pp. 13545–13551

Short Communications

Sunda Pangolin *Manis javanica* (Mammalia: Pholidota: Manidae) of Gaya Island, Sabah

– Jephte Sompud, Cynthia Boon Sompud, Kurtis Jai-Chyi Pei, Nick Ching-Min Sun, Rimi Repin & Fred Tuh, Pp. 13552–13556

Distribution and morphometric measurements of Blanford's Fox Vulpes cana (Mammalia: Carnivora: Canidae) of the Kingdom of Saudi Arabia

- Abdulhadi Aloufi & Ehab Eid, Pp. 13557-13562

Sebaceous gland adenoma in a free-ranging Baird's Tapir Tapirus bairdii (Tapiridae: Perissodactyla)

 Randall Arguedas, Maricruz Guevara-Soto & Jorge Rojas-Jiménez, Pp. 13563–13566

Recent records of the Banded Racer *Argyrogena fasciolata* (Shaw, 1802) (Reptilia: Squamata: Colubridae) from southern Coromandel Coast, peninsular India

Janani Sagadevan, Sumaithangi Rajagopalan Ganesh, Nitesh Anandan
 & Raveen Rajasingh, Pp. 13567–13572

A new species of *Simulium* (*Simulium*) (Diptera: Simuliidae), with keys to *S. striatum* species-group from India

- Sankarappan Anbalagan, Suryliyandi Vijayan, Chellapandian Balachandran & Sundaram Dinakaran, Pp. 13573–13578

New host records of polyphagous Lepidoptera on Ban Oak *Quercus leucotrichophora* A. Camus (Fabaceae) in the Garhwal Himalaya, India

 – Arun Pratap Singh, Kalpana Bahuguna & Gaurav Chand Ramola, Pp. 13579–13591

A preliminary study of the hawkmoth diversity (Lepidoptera: Sphingidae) of Kanyakumari District, Tamil Nadu, India – Geetha Iyer & Ian James Kitching, Pp. 13592–13604

Calamus pseudoerectus (Arecaceae), a new species from the eastern Himalaya, India

- Sujit Mondal, Shyamal K. Basu & Monoranjan Chowdhury, Pp. 13605-13610

Weed diversity in rice crop fields of Fatehgarh Sahib District, Punjab, India

Yadvinder Singh & Rai Singh, Pp. 13611–13616

Observations on the female flowers and fruiting of Tape Grass Enhalus acoroides from South Andaman Islands, India – Vardhan Patankar, Tanmay Wagh & ZoyaTyabji, Pp. 13617–13621

Notes

First records of *Agnidra vinacea* (Moore, 1879) (Lepidoptera: Drepanidae: Drepaninae) from the western Himalaya, extending its known range westwards Driba Day & Sanjay Sandhi, Dp. 12622, 12624

- Pritha Dey & Sanjay Sondhi, Pp. 13622-13624

Pollinators of Sikkim Mandarin Orange *Citrus reticulata* (Sapindales: Rutaceae)

– Urbashi Pradhan & M. Soubadra Devy, Pp. 13625–13628

Book Review

A holistic look on birds in urban areas

- S. Suresh Ramanan & Lalit Upadhyay, Pp. 13629-13630

Publisher & Host



Member



