

The Journal of Threatened Taxa (JoTT) is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at <a href="www.threatenedtaxa.org">www.threatenedtaxa.org</a>. All articles published in JoTT are registered under Creative Commons Attribution 4.0 International License unless otherwise mentioned. JoTT allows unrestricted use, reproduction, and distribution of articles in any medium by providing adequate credit to the author(s) and the source of publication.

# **Journal of Threatened Taxa**

Building evidence for conservation globally

www.threatenedtaxa.org

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

# **SHORT COMMUNICATION**

EURYLOPHELLA KARELICA TIENSUU, 1935 (INSECTA: EPHEMEROPTERA: EPHEMERELLIDAE) — AN ADDITIONAL SPECIES TO THE MAYFLY FAUNA OF UKRAINE AND NOTES ON DISTRIBUTION OF THE FAMILY IN THE COUNTRY



Alexander V. Martynov



For Focus, Scope, Aims, Policies, and Guidelines visit https://threatenedtaxa.org/index.php/JoTT/about/editorialPolicies#custom-0 For Article Submission Guidelines, visit https://threatenedtaxa.org/index.php/JoTT/about/submissions#onlineSubmissions For Policies against Scientific Misconduct, visit https://threatenedtaxa.org/index.php/JoTT/about/editorialPolicies#custom-2 For reprints, contact <ravi@threatenedtaxa.org>

The opinions expressed by the authors do not reflect the views of the Journal of Threatened Taxa, Wildlife Information Liaison Development Society, Zoo Outreach Organization, or any of the partners. The journal, the publisher, the host, and the partners are not responsible for the accuracy of the political boundaries shown in the maps by the authors.

Member



**Publisher & Host** 



# Journal of Threatened Taxa | www.threatenedtaxa.org | 26 April 2020 | 12(5): 15651-15654

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

DOI: https://doi.org/10.11609/jott.5692.12.5.15651-15654

#5692 | Received 09 January 2020 | Final received 18 February 2020 | Finally accepted 04 April 2020







# Eurylophella karelica Tiensuu, 1935 (Insecta: Ephemeroptera: Ephemerellidae) – an additional species to the mayfly fauna of Ukraine and notes on distribution of the family in the country

Alexander V. Martynov (1)

National Museum of Natural History, National Academy of Sciences of Ukraine, Bohdan Khmelnytsky str., 15, 01601 Kiev, Ukraine. martynov\_av@ukr.net, centroptilum@gmail.com

**Abstract**: A rare European species, *Eurylophella karelica* Tiensuu, 1935 (Ephemeroptera: Ephemerellidae) is reported for the first time from Ukraine. The larvae of the species were found within Ukrainian Polissya region in the Pripyat' River basin. Species list of family Ephemerellidae of Ukraine with notes on species distribution within the country is given in the present contribution.

**Keywords:** Checklist, distribution, mayfly, Pannota, species, Ukrainian Polissya region.

Eurylophella Tiensuu, 1935 is a Holarctic genus encompassing 19 recent species. Most of these species inhabit the Neartcic realm, and only three of them occur within the western Palearctic region—Eurylophella karelica Tiensuu, 1935, E. iberica Keffermüller & Da Terra, 1978, and E. korneyevi Martynov, Palatov & Godunko, 2015 (Martynov et al. 2015). Eurylophella karelica is the most common and widely distributed, the other two are rare with restricted distribution. Distribution of E. karelica extends from north to south, and falls within the north-west of the European part of Russia, Finland, Lithuania, Latvia, Estonia, Poland, Hungary, and Slovenia (Martynov et al. 2015; Ozoliņš et al. 2017). This species has not yet been recorded in Belarus (Moroz & Lipinskaya 2014), but it is likely to be found in the future.

# **MATERIALS AND METHODS**

The material was collected with square hydrobiological hand net 25cm long on each side. The sample is stored in 90–95 % ethanol in the author's collection in the National Museum of Natural History, National Academy of Sciences of Ukraine (Kyiv, Ukraine); its inventory number (IN) is Riv19Eurkar. Toponyms and hydronyms are duplicated in Ukrainian in square brackets.

Habitus of *E. karelica* larva was photographed using a Leica Z16 APO stereomicroscope equipped with Leica DFC450 Digital Camera in the I.I. Schmalhausen Institute of Zoology, National Academy of Sciences of Ukraine, and was subsequently processed with LAS Core 3.8 and Adobe® Photoshop™ CS5 software.

Material: Five larvae of Eurylophella karelica, Ukraine, Rivne region [Рівненська область], vicinity of Osnyts'k [Осницьк] Village, L'va [Льва] River, 51.284E, 27.145N and 51.284°E, 27.144°N (Image 1), about 145m, 22.xii.2018, leg. Martynov A.V. – IN Riv19Eurkar.

# **RESULTS AND DISCUSSION**

In Ukraine, *E. karelica* was recorded at Polissya [Полісся] region in potamal zone of L'va [Льва] River

Editor: K.A. Subramanian, Zoological Survey of India, Chennai, India.

Date of publication: 26 April 2020 (online & print)

Citation: Martynov, A.V. (2020). Eurylophella karelica Tiensuu, 1935 (Insecta: Ephemeroptera: Ephemerellidae) – an additional species to the mayfly fauna of Ukraine and notes on distribution of the family in the country. Journal of Threatened Taxa 12(5): 15651–15654. https://doi.org/10.11609/jott.5692.12.5.15651-15654

Copyright: © Martynov 2020. Creative Commons Attribution 4.0 International License. JoTT allows unrestricted use, reproduction, and distribution of this article in any medium by providing adequate credit to the author(s) and the source of publication.

Funding: IDEA WILD (some equipment used in this research)

Competing interests: The author declares no competing interests.

Acknowledgements: Author is grateful to Dr. M.V. Franchuk (Rivnenskyi Nature Reserve, Ukraine) and Dr. A.M. Roman (independent researcher) for the comprehensive help in the joint expeditions within Ukrainian Polissya. Some equipment used in this research was provided by IDEA WILD organization in 2017 and 2018.






Image 1. Geographical location of Eurylophella karelica Tiensuu, 1935 collecting place: a—Rivne Region | b—vicinity of Osnyts'k Village on map of Rivne Region | c—total view of L'va River watercourse in vicinity of Osnyts'k Village.

(Image 2) that belongs to Pripyat' [Прип'ять] River basin. This finding is the first record of *E. karelica* from Ukraine. The Pripyat' River basin, where the species was found, differs in the presence of a relatively large number of waterbodies, with slightly impaired species compositions of aquatic insects, in comparison with vast parts of lowland Ukraine. Thus, during investigation of Ephemeroptera at the Pripyat' River basin in 2017–2018 several stenobiotic and rare species were registered (Martynov 2018); some of these species had not yet been recorded elsewhere in Ukraine.

# **Habitat characteristics**

The section of the L'va River, where *E. karelica* larvae were collected (Image 2, 3), is situated downstream of a dam. The river has moderate current velocity (up to 0.3m/s), but under the bridge and above the dam rapids with strong current are present (up to 1m/s). The bottom of the river is silty, sandy (in some places silted), and stony (under the bridge). All larvae of the species were collected from dead aquatic vegetation (leaves, stems etc.) in small bays with silted bottom and almost lacking flow (Image 2). The additional parameters of waterbody in collecting sites measured during the sampling (22. xii.2018) were: water temperature 6°C, water hardness 84ppm, pH 8.9.

The following mayflies were also collected within the investigated section of the L'va River in winter (22. xii.2018) and spring (22.iii.2019)—*Nigrobaetis digitatus* 

(Bengtsson 1912), Cloeon dipterum (Linnaeus, 1761), Heptagenia sulphurea (O.F. Müller, 1776), H. flava Rostock, 1878, Kageronia fuscogrisea (Retzius, 1783), Leptophlebia marginata (Linnaeus, 1767), L. vespertina (Linnaeus, 1758), Paraleptophlebia submarginata (Stephens, 1835), Ephemera vulgata Linnaeus, 1758, and Caenis sp. The registration of 11 mayfly species in this section of the L'va River, makes me consider it as one of the hotspots of mayfly species diversity in Ukrainian Polissya region.

# Ephemerellidae of Ukraine and their distribution within country

In the first checklist of mayflies of Ukraine, five species of subfamily Ephemerellinae were mentioned (Godunko & Kłonowska-Olejnik 2003). *Eurylophella karelica,* recorded for Ukraine for the first time herein, belongs to another subfamily — Timpanoginae. Thereby, now species list of Ephemerellidae of Ukraine consists of six species from five genera and two subfamilies, and *E. karelica* is the most rare species among them.

# **Ephemerellinae**

- · Serratella ignita (Poda, 1761)
- Ephemerella notata Eaton, 1887
- · Ephemerella mucronata (Bengtsson, 1909)
- · Teloganopsis mesoleuca (Brauer, 1857)
- · Torleya major (Klapálek, 1905)





Image 2–3. Larva of Eurylophella karelica Tiensuu, 1935 and its habitat: 2—total view of the L'va River in collecting cite (area outlined with white color - microhabitat mostly preferred by E. karelica) | 3—larva of E. karelica, dorsal view. Scale bar 2mm. © Alexander V. Martynov.

# **Timpanoginae**

· Eurylophella karelica Tiensuu, 1935 – first record for Ukraine

The Carpathian Mountains [Карпати], Transcarpathian [Закарпатська] Lowland, eastern Carpathian Foothills [Передкарпаття] are the Ukrainian regions inhabited with the highest number of Ephemerellidae taxa; a total of five species are recorded here—S. ignita, E. notata, E. mucronata, T. mesoleuca, and T. major (Mikulski 1935; Godunko 2000; Godunko & Kłonowska-Olejnik 2003; Afanasyev 2006; Kovács & Godunko 2008). Such diversity is reasoned by abundance of waterbodies with unpolluted or moderately polluted rhithral and epipotamal zones preferred by a vast

number of mayfly species.

Except for *E. karelica*, also *S. ignita* inhabits some waterbodies of Ukrainian Polissya. Larvae of this last species were registered in Dnipro [Дніпро] River (Zimbalevskaya et al. 1989), Pripyat' River (Trylis et al. 2013), and Irsha [Ірша] River (50.759E & 29.411N) (original data). Such small numbers of species in the region may be due to the small number of flowing waterbodies preferred by Ephemerellidae, pollution and habitat modifications in main part of them.

Four species of Ephemerellidae, Serratella ignita, Teloganopsis mesoleuca, Ephemerella notata, and E. mucronata were recorded for Dnister [Дністер] River, natural border of Podolia [Подільська] Upland, and



eastern Carpathian Foothills (Telyuk 1982, 1992; Mikulski 1933).

Mayflies of Podolia and Volyn' [Волинська] Uplands, lowland, central, and southern regions of Ukraine are still poorly investigated. Most of the data are old and need confirmation due to significant changes and pollution levels in waterbodies, especially in central and southern regions.

The Crimean [Кримські] Mountains was investigated and summarized in detail (see overview in Prokopov & Godunko 2007). Only *Serratella ignita* was listed by Kiseleva & Yezernitskiy (1985) for this region, and there are no other mentions of the species for the Crimea [Крим]. According to Prokopov & Godunko (2007) the presence of this species in the fauna of the region needs confirmation.

Within physiographic regions of eastern Ukraine only one representative of Ephemerellidae—Serratella ignita—was registered. This species inhabits only rhithral and eupotamal zones of streams and headwaters of rivers within most elevated parts of Donetsk [Донецька] elevated areas, and the species is rare within it (Martynov 2014, 2016). The recorded population of *S. ignita* is geographically isolated and should be considered as glacial relicts in this territory.

Itshould be noted, that the diversity of Ephemerellidae in neighboring countries (Romania, Hungary, Slovakia, Poland, and European part of Russian Federation) does not exceed diversity of the family within Ukraine. The territory of Moldova is poorly investigated; two species of the family were recorded for Belarus only. Therefore, the Ephemerellidae species list given above should be considered as complete, and no other species of the family are expected to be registered in Ukraine in the near future.

# REFERENCES

- Afanasyev, S.A. (2006). Biotic communities structure and assessment of Tisa's basin rivers ecological status. Kyiv, Interdruk, 101pp. https://doi.org/10.13140/RG.2.1.2864.0160 (in Ukrainian, English summary)
- **Godunko, R.J. (2000).** Historycal changes of the fauna and questions of reservation of mayfly (Ephemeroptera, Insecta) of Ukrainian Carpathians. *Proceedings of the State Natural History Museum* 15: 158–168. (in Ukrainian, English summary).
- Godunko, R.J. & M. Kłonowska-Olejnik (2003). A checklist of the Ukrainian mayflies (Insecta: Ephemeroptera). *Polskie Pismo Entomologiczne* 72: 203–210.

- Trylis, V.V., O.E. Usov, O.M. Lietytska, S.O. Afanasyev, V.A. Martinchuk
  T.V. Oberemchuk (2013). Macroinvertebretes. pp: 163–175.
  In: Romanenko, V.D., S.O. Afanasyev and V.I. Osadchiy (eds.).
  Hydroecosystems of the protected territories of the upper Prypiat' under climate changes: monograph. Kyiv, Kafedra, 228pp. (in Ukrainian)
- Kiseleva, G.A. & E.V. Yezernitskiy (1985). Spatial distribution of aquatic entomofauna in the upper part of Salgir River basin under anthropogenic influence. *Ecological and environmental aspects of the study of the Crimean Mountains*. Simferopol: SGU: 110–119. (in Russian)
- Kovács, T. & R.J. Godunko (2008). Faunistical records of larvae of Ephemeroptera, Odonata and Plecoptera from the Zakarpats'ka Region, Ukraine. Folia Historico Naturalia Musei Matraensis 32: 87–91.
- Martynov, A.V. (2014). Mayflies (Ephemeroptera) of Rhithral and Crenal Zones of Donetsk Elevated Physiographic Area's Watercourses (Eastern Ukraine): species composition, ecological characteristics. Caucasian Entomological Bulletin 10(1): 3–18. (in Russian, English summary)
- Martynov, A.V. (2016). The Life Cycles of Mayflies (Insecta: Ephemeroptera) of the Eastern Ukraine. Second Report. *Proceedings of the National Museum of Natural History* 14: 86–94.
- Martynov, A.V. (2018). New records of some rare mayflies (Insecta, Ephemeroptera) from Ukraine. *Ecologica Montenegrina* 16: 48–57.
- Martynov, A.V., D.M. Palatov & R.J. Godunko (2015). The larvae of West Palaearctic *Eurylophella* Tiensuu, 1935 (Ephemeroptera: Ephemerellidae), with description of a new species from Georgia. *Zootaxa* 3904(1): 123–143. https://doi.org/10.11646/zootaxa.3904.1.8
- Mikulski, J. (1933). Próba charakterystyki zoogeograficznej jętek (Ephemeroptera) fauny Polski. *Pamietnik XIV Zjazdu Lekarzy i Przyrodników Polskich w Poznaniu*: 476–478.
- Mikulski, J.S. (1935). Jętki (Ephemeroptera), pp. 63–71. In: *Przyczynek do znajomości fauny Czarnohory*. Rozprawy i sprawozdanie. Instytut Badawczy lasów państwowych. Serjia A, 8, Warszawa, Pologne: 27–102 pp.
- Moroz, M.D. & T.P. Lipinskaya (2014). A Cataloge of mayflies (Ephemeroptera), stoneflies (Plecoptera) and caddisflies (Trichoptera) of Belarus. Bielaruskaja navuka, Minsk, 315pp. (in Russian)
- Ozoliņš, D., J. Jēkabsone, A. Skuja & H. Timm (2017). New records of a poorly studied mayfly species, *Eurylophella karelica* (Ephemeroptera, Ephemerellidae) Tiensuu, 1935, in the Baltic Ecoregion. *Check List* 13(4): 349–353. https://doi.org/10.15560/13.4.349
- Prokopov, G.A. & R.J. Godunko (2007). A Catalogue of the mayflies (Insecta: Ephemeroptera) of the Crimea. Proceeding of IV International Scientific and Practical Conference. Part 2, Zoology: 140–152pp. (in Russian)
- **Telyuk, P.M. (1982).** Zoobenthos of Dniester River. PhD Thesis Abstract. Institute of Hydrobiology of the UkrSSR, Kyiv, 16pp. (in Russian)
- Telyuk, P.M. (1992). Macrozoobenthos of upper part of Dniester River, pp. 245–249. In: Braginsky, L.P. (ed.). *Hydrobiological regime of Dniester River and its waterbodies*. Kyiv, Naukova dumka, 356pp. (in Russian)
- Zimbalevskaya, L.N., P.G. Sukhoivan, M.I. Chernogorenko et al. (1989). Invertebrates and fishes of the Dnieper and its reservoirs. Kyiv, Naukova dumka, 247pp. (in Russian)





The Journal of Threatened Taxa (JoTT) is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at www.threatenedtaxa.org. All articles published in JoTT are registered under Creative Commons Attribution 4.0 International License unless otherwise mentioned. JoTT allows allows unrestricted use, reproduction, and distribution of articles in any medium by providing adequate credit to the author(s) and the source of publication.

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

April 2020 | Vol. 12 | No. 5 | Pages: 15535–15674 Date of Publication: 26 April 2020 (Online & Print) DOI: 10.11609/jott.2020.12.5.15535-15674

www.threatenedtaxa.org

#### Article

Prey selection and food habits of the Tiger *Panthera tigris* (Mammalia: Carnivora: Felidae) in Kalakkad-Mundanthurai Tiger Reserve, southern Western Ghats, India

Bawa Mothilal Krishnakumar, Rajarathinavelu Nagarajan & Kanagaraj Muthamizh Selvan, Pp. 15535–15546

## **Communications**

Community-based study to demonstrate the presence and local perspectives of the Critically Endangered Chinese Pangolin *Manis pentadactyla* in Zhejiang Wuyanling, China

Hongying Li, Shusheng Zhang, Ji Zhang, Zupei Lei, Fangdong Zheng
 Peter Daszak, Pp. 15547–15556

## Field friendly method for wild feline semen cryopreservation

 Gediendson Ribeiro de Araujo, Thyara de Deco-Souza, Letícia Coelho Ferreira Bergo, Leanes Cruz da Silva, Ronaldo Gonçalves Morato, Pedro Nacib Jorge-Neto, Maitê Cardoso Coelho da Silva, Gustavo Guerino Macedo & Tarcízio Antônio Rego De Paula, Pp. 15557–15564

Habitat structure determines the abundance of the Endangered Sharpe's Longclaw *Macronyx sharpei* (Aves: Passeriformes: Motacillidae) at Timau montane grasslands in central Kenya

 Dominic Kimani, Muchane Muchai, Johnstone Kimanzi, Joseph Mwangi, Wanyoike Wamiti, Samuel Bakari, Bernhard Walter & Peter Njoroge, Pp. 15565–15571

Avifaunal diversity of some selected water bodies of Khanapur Taluka, Belagavi District, Karnataka, India

Harsha D. Neelgund & Girish Kadadevaru, Pp. 15572–15586

Herpetofauna of Shuklaphanta National Park, Nepal

Yam Bahadur Rawat, Santosh Bhattarai, Laxman Prasad Poudyal & Naresh Subedi, Pp. 15587–15611

Varying colour pattern, yet genetically similar: Pebble Crab Seulocia vittata (Stimpson, 1858) (Brachyura: Leucosiidae) from the southeastern coast of India

– Sanjeevi Prakash & Amit Kumar, Pp. 15612–15618

Grasses of Kundadri Hills in the Western Ghats of Karnataka, India

Hanchali Udayashankar Abhijit & Yelugere Linganaik Krishnamurthy,
 Pp. 15619–15630

# Comparative phytosociological assessment of three terrestrial ecosystems of Wayanad Wildlife Sanctuary, Kerala, India

– M. Vishnu Chandran, S. Gopakumar & Anoopa Mathews, Pp. 15631–15645

### **Short Communications**

Piroplasmosis in a captive Grant's Zebra *Equus quagga boehmi* (Mammalia: Perissodactyla: Equidae) - a case study

 Sarat Kumar Sahu, Niranjana Sahoo, Bijayendranath Mohanty & Debabrat Mohapatra, Pp. 15646–15650

Eurylophella karelica Tiensuu, 1935 (Insecta: Ephemeroptera: Ephemerellidae) – an additional species to the mayfly fauna of Ukraine and notes on distribution of the family in the country

- Alexander V. Martynov, Pp. 15651-15654

Some new records of katydids (Orthoptera: Tettigoniidae) from Uttar Pradesh, India

- Ramesh Singh Yadav & Dharmendra Kumar, Pp. 15655-15660

# Notes

On the occurrence of Honey Badger *Mellivora capensis* (Mammalia: Carnivora: Mustelidae) in the northern Eastern Ghats of Andhra Pradesh, India

Vikram Aditya, Yogesh Pasul & Ganesh Thyagarajan, Pp. 15661– 15663

Assamese Cat Snake *Boiga quincunciata* (Wall, 1908) (Reptilia: Squamata: Colubridae) - new country record for Bhutan

– Lekey Chaida, Abhijit Das, Ugyen Tshering & Dorji Wangdi,Pp. 15664–15667

Loss of Critically Endangered Hawksbill Turtle nesting beach at EGA facility, Abu Dhabi, UAE

– D. Adhavan, Pp. 15668–15670

*Meliola elaeocarpicola* sp. nov. (Ascomycetes, Meliolales) from Malabar Wildlife Sanctuary in Kerala State, India

- Lini K. Mathew & Jacob Thomas, Pp. 15671-15674

**Publisher & Host** 





