


Building evidence for conservation globally

Journal of Threatened TAXA



Open Access

10.11609/jott.2024.16.7.25495-25638
www.threatenedtaxa.org

26 July 2024 (Online & Print)
16 (7): 25495-25638
ISSN 0974-7907 (Online)
ISSN 0974-7893 (Print)

- Lakshmi Niranjana -



Publisher

Wildlife Information Liaison Development Societywww.wild.zooreach.org

Host

Zoo Outreach Organizationwww.zooreach.org43/2 Varadarajulu Nagar, 5th Street West, Ganapathy, Coimbatore, Tamil Nadu 641006, India

Registered Office: 3A2 Varadarajulu Nagar, FCI Road, Ganapathy, Coimbatore, Tamil Nadu 641006, India

Ph: +91 9385339863 | www.threatenedtaxa.orgEmail: sanjay@threatenedtaxa.org**EDITORS****Founder & Chief Editor****Dr. Sanjay Molur**Wildlife Information Liaison Development (WILD) Society & Zoo Outreach Organization (ZOO),
43/2 Varadarajulu Nagar, 5th Street West, Ganapathy, Coimbatore, Tamil Nadu 641006, India**Deputy Chief Editor****Dr. Neelesh Dahanukar**

Noida, Uttar Pradesh, India

Managing Editor**Mr. B. Ravichandran**, WILD/ZOO, Coimbatore, Tamil Nadu 641006, India**Associate Editors****Dr. Mandar Paingankar**, Government Science College Gadchiroli, Maharashtra 442605, India**Dr. Ulrike Streicher**, Wildlife Veterinarian, Eugene, Oregon, USA**Ms. Priyanka Iyer**, ZOO/WILD, Coimbatore, Tamil Nadu 641006, India**Dr. B.A. Daniel**, ZOO/WILD, Coimbatore, Tamil Nadu 641006, India**Editorial Board****Dr. Russel Mittermeier**

Executive Vice Chair, Conservation International, Arlington, Virginia 22202, USA

Prof. Mewa Singh Ph.D., FASc, FNA, FNAsc, FNAPsy

Ramanna Fellow and Life-Long Distinguished Professor, Biopsychology Laboratory, and Institute of Excellence, University of Mysore, Mysuru, Karnataka 570006, India; Honorary Professor, Jawaharlal Nehru Centre for Advanced Scientific Research, Bangalore; and Adjunct Professor, National Institute of Advanced Studies, Bangalore

Stephen D. Nash

Scientific Illustrator, Conservation International, Dept. of Anatomical Sciences, Health Sciences Center, T-8, Room 045, Stony Brook University, Stony Brook, NY 11794-8081, USA

Dr. Fred Pluthero

Toronto, Canada

Dr. Priya Davidar

Sigur Nature Trust, Chadapatti, Mavinahalli PO, Nilgiris, Tamil Nadu 643223, India

Dr. Martin Fisher

Senior Associate Professor, Battcock Centre for Experimental Astrophysics, Cavendish Laboratory, JJ Thomson Avenue, Cambridge CB3 0HE, UK

Dr. John Fellowes

Honorary Assistant Professor, The Kadoorie Institute, 8/F, T.T. Tsui Building, The University of Hong Kong, Pokfulam Road, Hong Kong

Prof. Dr. Mirco Solé

Universidade Estadual de Santa Cruz, Departamento de Ciências Biológicas, Vice-coordenador do Programa de Pós-Graduação em Zoologia, Rodovia Ilhéus/Itabuna, Km 16 (45662-000) Salobrinho, Ilhéus - Bahia - Brasil

Dr. Rajeev Raghavan

Professor of Taxonomy, Kerala University of Fisheries & Ocean Studies, Kochi, Kerala, India

English Editors**Mrs. Mira Bhojwani**, Pune, India**Dr. Fred Pluthero**, Toronto, Canada**Mr. P. Ilangovan**, Chennai, India**Ms. Sindhura Stothra Bhashyam**, Hyderabad, India**Web Development****Mrs. Latha G. Ravikumar**, ZOO/WILD, Coimbatore, India**Typesetting****Mrs. Radhika**, ZOO, Coimbatore, India**Mrs. Geetha**, ZOO, Coimbatore India**Fundraising/Communications****Mrs. Payal B. Molur**, Coimbatore, India**Subject Editors 2020–2022****Fungi****Dr. B. Shivaraju**, Bengaluru, Karnataka, India**Dr. R.K. Verma**, Tropical Forest Research Institute, Jabalpur, India**Dr. Vatsavaya S. Raju**, Kakatiya University, Warangal, Andhra Pradesh, India**Dr. M. Krishnappa**, Jnana Sahyadri, Kuvenpu University, Shimoga, Karnataka, India**Dr. K.R. Sridhar**, Mangalore University, Mangalagangotri, Mangalore, Karnataka, India**Dr. Gunjan Biswas**, Vidyasagar University, Midnapore, West Bengal, India**Dr. Kiran Ramchandra Ranadive**, Annasaheb Magar Mahavidyalaya, Maharashtra, India**Plants****Dr. G.P. Sinha**, Botanical Survey of India, Allahabad, India**Dr. N.P. Balakrishnan**, Ret. Joint Director, BSI, Coimbatore, India**Dr. Shonil Bhagwat**, Open University and University of Oxford, UK**Prof. D.J. Bhat**, Retd. Professor, Goa University, Goa, India**Dr. Ferdinand Boero**, Università del Salento, Lecce, Italy**Dr. Dale R. Calder**, Royal Ontario Museum, Toronto, Ontario, Canada**Dr. Cleofas Cervancia**, Univ. of Philippines Los Baños College Laguna, Philippines**Dr. F.B. Vincent Florens**, University of Mauritius, Mauritius**Dr. Merlin Franco**, Curtin University, Malaysia**Dr. V. Irudayaraj**, St. Xavier's College, Palayamkottai, Tamil Nadu, India**Dr. B.S. Kholia**, Botanical Survey of India, Gangtok, Sikkim, India**Dr. Pankaj Kumar**, Department of Plant and Soil Science, Texas Tech University, Lubbock, Texas, USA**Dr. V. Sampath Kumar**, Botanical Survey of India, Howrah, West Bengal, India**Dr. A.J. Solomon Raju**, Andhra University, Visakhapatnam, India**Dr. Vijayasankar Raman**, University of Mississippi, USA**Dr. B. Ravi Prasad Rao**, Sri Krishnadevaraya University, Anantapur, India**Dr. K. Ravikumar**, FRLHT, Bengaluru, Karnataka, India**Dr. Aparna Watve**, Pune, Maharashtra, India**Dr. Qiang Liu**, Xishuangbanna Tropical Botanical Garden, Yunnan, China**Dr. Noor Azhar Mohamed Shazili**, Universiti Malaysia Terengganu, Kuala Terengganu, Malaysia**Dr. M.K. Vasudeva Rao**, Shiv Ranjan Housing Society, Pune, Maharashtra, India**Prof. A.J. Solomon Raju**, Andhra University, Visakhapatnam, India**Dr. Mander Datar**, Agharkar Research Institute, Pune, Maharashtra, India**Dr. M.K. Janarthanam**, Goa University, Goa, India**Dr. K. Karthigeyan**, Botanical Survey of India, India**Dr. Errol Vela**, University of Montpellier, Montpellier, France**Dr. P. Lakshminarasiham**, Botanical Survey of India, Howrah, India**Dr. Larry R. Noblick**, Montgomery Botanical Center, Miami, USA**Dr. K. Haridasan**, Pallavur, Palakkad District, Kerala, India**Dr. Analinda Manila-Fajard**, University of the Philippines Los Baños, Laguna, Philippines**Dr. P.A. Siru**, Central University of Kerala, Kasaragod, Kerala, India**Dr. Afroz Alam**, Banasthali Vidyapith (accredited A grade by NAAC), Rajasthan, India**Dr. K.P. Rajesh**, Zamorin's Guruvayurappan College, GA College PO, Kozhikode, Kerala, India**Dr. David E. Boufford**, Harvard University Herbaria, Cambridge, MA 02138-2020, USA**Dr. Ritesh Kumar Choudhary**, Agharkar Research Institute, Pune, Maharashtra, India**Dr. A.G. Pandurangan**, Thiruvananthapuram, Kerala, India**Dr. Navendu Page**, Wildlife Institute of India, Chandrabani, Dehradun, Uttarakhand, India**Dr. Kannan C.S. Warrier**, Institute of Forest Genetics and Tree Breeding, Tamil Nadu, India**Invertebrates****Dr. R.K. Avasthi**, Rohtak University, Haryana, India**Dr. D.B. Bastawade**, Maharashtra, India**Dr. Partha Pratim Bhattacharjee**, Tripura University, Suryamaninagar, India**Dr. Kailash Chandra**, Zoological Survey of India, Jabalpur, Madhya Pradesh, India**Dr. Ansie Dippenaar-Schoeman**, University of Pretoria, Queenswood, South Africa**Dr. Rory Dow**, National Museum of natural History Naturalis, The Netherlands**Dr. Brian Fisher**, California Academy of Sciences, USA**Dr. Richard Gallon**, Ilandudno, North Wales, LL30 1UP**Dr. Hemanth V. Ghate**, Modern College, Pune, India**Dr. M. Monwar Hossain**, Jahangirnagar University, Dhaka, BangladeshFor Focus, Scope, Aims, and Policies, visit https://threatenedtaxa.org/index.php/JoTT/aims_scopeFor Article Submission Guidelines, visit <https://threatenedtaxa.org/index.php/JoTT/about/submissions>For Policies against Scientific Misconduct, visit https://threatenedtaxa.org/index.php/JoTT/policies_various

continued on the back inside cover

Cover: Mixed media illustration of a Blue bird and Sunbird. © Lakshmi Niranjan.



***Heleocoris stephanus* (Heteroptera: Naucoridae: Laccocorinae), a new species of creeping water bug from Kallada River, Kerala, India**

Dani Benchamin¹ , R. Sreejai²  & M.S. Arya³ 

¹⁻³ Zoology Research Centre, St. Stephen's College, Pathanapuram, University of Kerala, Thiruvananthapuram, Kerala 695034, India.

¹danibenchamin93@gmail.com (corresponding author), ²sreejaiksbb@gmail.com, ³aryamadhussc@gmail.com

Abstract: A new species of *Heleocoris*, Stål (1876) was collected from the upstream Kallada River in Kerala, southwestern India. This discovery brings the number of described species for this genus in India to twelve. This species can be diagnosed by the overall body coloration and distinct morphological & genital features. Descriptions including a habitus photograph, ecology, and diagnostic comparisons among congeners are provided.

Keywords: Coloration, congeners, genitalia, morphology, upstream, Urukunnu.

Malayalam: പൊലിഡാകോറിസ് ഇന്ത്യൻ പെട്ട ഒരു പുതിയ സ്പീഷിസിനെ ഇന്ത്യയുടെ തെക്കുപടിഞ്ഞാറൻ സംസ്ഥാനമായ കേരളത്തിലെ കല്ലാട നദിയിൽ നിന്ന് കണ്ടെത്തി. ഈ കണ്ടെത്തലോടു കൂടി ഇന്ത്യയിൽ ഈ ഇന്ത്യൻ സ്പീഷിസുകളുടെ ഏറ്റവും പാരിശോധി. ശർഭത്തിലെ വ്യത്യസ്തതിയിലൂടെ ഒക്കമായ നിറവും അഭ്യാസങ്ങളും രൂപാംഗത്പരവും ജനങ്ങളുടെ സംസ്ഥാനമായ മറ്റു സവിശേഷകളുമാണ് ഈ പുതിയ സ്പീഷിസിനെ മറ്റു സ്പീഷിസുകളിൽ നിന്നു വ്യത്യസ്തമാക്കുന്നത്. വിശദമായ വിവരങ്ങളും അനുബന്ധ ചിത്രങ്ങളും താരതമ്യ പാനങ്ങൾ ഉൾപ്പെടെയുള്ള വിവരങ്ങളും ഈ പ്രസിദ്ധീകരണത്തിൽ നൽകിയിരിക്കുന്നു.

ZooBank: urn:lsid:zoobank.org:pub:8F58793E-3BD5-4EB6-A07B-3DB1BF7A63D7

Editor: Hasko Nesemann, Central University of Bihar, Patna, India.

Date of publication: 26 July 2024 (online & print)

Citation: Benchamin, D., R. Sreejai & M.S. Arya (2024). *Heleocoris stephanus* (Heteroptera: Naucoridae: Laccocorinae), a new species of creeping water bug from Kallada River, Kerala, India. *Journal of Threatened Taxa* 16(7): 25584-25589. <https://doi.org/10.11609/jott.9129.16.7.25584-25589>

Copyright: © Benchamin et al. 2024. Creative Commons Attribution 4.0 International License. JoTT allows unrestricted use, reproduction, and distribution of this article in any medium by providing adequate credit to the author(s) and the source of publication.

Funding: Self-funded.

Competing interests: The authors declare no competing interests.

Author details: DR. DANI BENCHAMIN, a former PhD scholar, Postgraduate and Research Department of Zoology, St. Stephen's College has research interests in freshwater ecology and biomonitoring. SREEJAI R serves as an assistant professor and research guide at the Postgraduate and Research Department of Zoology, St. Stephen's College, Pathanapuram, and DR. ARYA MS, a former PhD scholar of the same department has a research interest in estuarine ecology and planktonology.

Author contributions: DB and AMS carried out fieldwork, identification and manuscript preparation, SR reviewed the manuscript.

Acknowledgements: All glory, honor and praise to my Lord and Savior Jesus Christ. We are grateful to the authorities of St. Stephen's College, Pathanapuram, University of Kerala, for providing laboratory facilities for this discovery. We extend our gratitude to our friend, Sreenish S., for his assistance in the field. We remember our parents and family members for their prayers and love. The research stipend for first and third authors has been provided by the University of Kerala.



INTRODUCTION

The creeping water bugs are subaquatic, but some species are present in the water covering a rock surface (hygropetric). Most Indian species present adjacent to the streams. The genus *Heleocoris* Stål (1876) was first included in the division Laccocoraria, which later was raised to the subfamily Laccocorinae, Montandon (1897). *Heleocoris* contains 29 species, of which 11 are known from India (Polhemus & Polhemus 2013). The recent decade witnessed the discovery of a new species: *Heleocoris mcphersoni*, Sites and Vitheepradit, 2011 from Thailand. Globally, the Naucoridae consists of 401 species while the Indian sub-region consists of 20 species under seven genera (Chandra et al. 2017). There are few taxonomic records of *Heleocoris* in the southern part of India, and faunal surveys in waterfalls, streams of mountain ranges, and forests are still required. *Heleocoris stephanus* sp. nov. from Kollam District of Kerala state, is described as new to science in the present paper. This species was generally found associated with plant debris on the edge of the rivers. A diagnosis and comparison with congeners in Kerala, Tamil Nadu and other states congeners is presented..

MATERIALS AND METHODS

Bug samples were collected by hand-picking with either fingers or soft forceps and use of an aquatic net

to sweep through submerged vegetation and organic debris along shorelines of river Urukunnu, Kerala (Image 1). Specimens were placed into vials with 90% ethyl alcohol and brought to the laboratory of Zoology Research Centre, St. Stephen's College, Pathanapuram. Morphological terminology in the description mostly follows that of Xie & Liu (2015) and Sites & Vitheepradit (2011). All measurements are given in mm. Olympus CX33 and Weswox SZM-105 microscopes were used to obtain images and followed by its preparation with Photoshop CS5 (Adobe Systems Inc., San Jose, CA). The holotype (ZSI/WGRC/I.R.-INV.26976) was deposited in the Zoological Survey of India, Western Ghats Regional Centre (WGRC), Kozhikode, Kerala, India. Paratypes (SSCDZ/Hem01/2024 and SSCDZ/Hem02/2024) were deposited in the Department Museum, Zoology Research Centre, St. Stephen's College, Pathanapuram.

RESULTS

Heleocoris stephanus sp. nov. (Image 2)

urn:lsid:zoobank.org:act:D01E304C-38E9-412A-A296-09DFEB97BE86

Material examined: Holotype (ZSI/WGRC/I.R.-INV.26976): 07 October 2023, macropterous male, India: Urukunnu station, Kollam District, Kerala, 8.5905°N & 77.0124°E, 45 m, upstream Kallada River, coll. Dani & Arya, Paratypes (SSCDZ/Hem01/2024 and SSCDZ/



Image 1. Urukunnu Station in the upstream of Kallada River, Kollam District, Kerala, India, the type locality of *Heleocoris stephanus* sp. nov.




Image 2. Diagnostic features of *Heleocoris stephanus* sp. nov. (holotype): A—dorsal habitus | B—ventral habitus | C—phallotheca. © Dani Benchamin.

Hem02/2024): same data as primary type.

Description: Macropterous male. Holotype, length 9.78; width 6.20. Paratypes ($n = 4$), length 9.44–9.88 (mean = 9.69); maximum width 6.09–6.29 (mean = 6.17). General shape is ovate, slightly flattened; widest across embolia. Description of *Heleocoris stephanus* sp. nov. and other congeners are presented in Table 1.

Coloration: Overall body color is anteriorly yellowish with black mottling, scutellum dark brown with irregular yellow apex, yellowish, and thick median longitudinal patch. Head with black spotting, generally over the entire yellowish dorsal surface; patterns coalescent at the posterior margin, reducing anteriorly as black spots adjoining anterior margin, eyes blackish with

pale yellowish anterolateral margin, rostrum last two segments dark brownish. Pronotum with black markings scattered over the entire dorsal surface; laterally reduced to a few black spots; black coalescent patches in the mid-anterior region; a yellowish transverse band with irregular black blotches at the posterior margin. Hemelytra black. Forelegs and hindlegs are dorsally brownish and ventrally pastel yellowish. Connexiva yellowish-brown antero-posterior, showing checkered appearance (Image 2B). Ventrally yellowish cephalothorax, brownish abdomen, and pygophore (Image 2B).

Head: Densely punctured with black. Single sclerite at posterior corner of eye, directed ventrad, anterior

end simple. Dome-shaped, blackish spot on the base of the head in between the eyes. Compound eyes blackish, convergent anteriorly, convex on lateral posterior margin. Broad semi-circular labrum, rostrum short with three visible segments, first segment partially covered with labrum, each following segment progressively narrower, reaching to near anterior prothoracic coxae. Antennae, thickened and hairy: not extending to lateral margin of head.

Thorax: Pronotum broad and maximum width at rounded posterolateral corners; tuft of setae in anterior end, transverse sulcus band ending at anterior end of clavus with ventral tuft of setae; ventro-lateral margins with blackish punctations; tuft of elongate sensory setae on anterior surface behind eye. Scutellum is finely granulose, tumescent, and elevated above level of wings; sinuated lateral margins. Well-developed hindwings. Hemelytron blackish-brown; largely yellowish embolium with black spots.

Legs: Foreleg with large femur, strongly widened antero-basally, lateral margins with thick bunch of setae; tibia narrow, with bunch of setae at lateral margins; tarsus bristled and two-segmented with two razor shaped claws. Hindleg with wide femur have small bulbous projections at its posterior end, tibia narrow with a profusion of swimming hairs; tarsus with small dorsal bristles and ventral elongated tuft of setae.

Abdomen: Abdomen ventrally covered with appressed hair; large blackish spots; ventrally with heavy marginal spines on laterotergites III–VI and elongate light-coloured hair.

Genitalia: Pygophore brownish and hairy, posterior margin acuminate with dense brush; parameres reduced, blackish, and symmetrical; phallosoma elongate, linear, widest & rounded triangular at apex (Image 2C).

Female: Unknown

Diagnosis: *Heleocoris stephanus* sp. nov. is similar to the congener *H. vicinus* (Montandon, 1910), by size, and overall coloration. However, *H. stephanus* is notably different from *H. vicinus* by the finely granulose, dark brown scutellum with irregular yellow apex and thick median longitudinal yellowish patch. *H. majusculus* (Montandon, 1908) and *H. rotundatus* (Montandon, 1908) are other congeners reported from the neighbouring state of Tamil Nadu. The former drastically differs from *H. stephanus* by its entirely black scutellum and overall body size, while the latter differs from *H. stephanus* by its black coloured body beneath with greyish pilose and body size. *H. breviceps* (Montandon, 1897) is another congener reported from the Indian

states such as Andhra Pradesh, Himachal Pradesh, Maharashtra, and Tamil Nadu. This congener differs from *H. stephanus* by largely blackish abdomen beneath and brownish scutellum. As the name implies, *H. elongatus* (Montandon, 1897) differs from *H. stephanus* by its body shape. Additionally, *H. elongatus* can be distinguished from *H. stephanus* by its entirely yellowish body beneath. *H. elongatus* was reported from the Indian states Bihar, Madhya Pradesh, Maharashtra, and West Bengal. *H. acutus* (Spinola, 1837) was reported only from Maharashtra and it can be distinguished from *H. stephanus* by its distinct acute posterior angles of pronotum. *H. bengalensis* (Montandon, 1910) was reported from the central states such as Madhya Pradesh, Uttar Pradesh, and West Bengal. *H. stephanus* can be distinguished from this congener by the distinct characteristics of genitalia such as brownish & hairy pygophore, posterior margin acuminate with dense brush, and blackish & reduced parameres. *H. bergrothi* (Montandon, 1897), reported from the southern Indian states of Karnataka, Kerala, and Tamil Nadu is similar to *H. stephanus* in a few characteristics such as body size, shape, and v-shaped discal spot in scutellum. *H. stephanus* differs from *H. bergrothi* by its distinct colouration of scutellum, such as, dark brown with irregular yellow apex, thick & yellowish median longitudinal patch, and distinct black mottling of head & pronotum. Lastly, the closest congener *H. indicus* (Montandon, 1897) shares a few similar features with *H. stephanus* like flavescent with brown punctures on head and pronotum, overall coloration, and v-shaped discal spot in scutellum. *H. indicus* has a number of contrasting characteristics such as entirely yellowish underbody, pronotum with blackish transverse colouration on the posterior margin, while the *H. stephanus* has a ventrally yellowish cephalothorax, brownish abdomen and pygophore, pronotum with irregular black blotches on the posterior transverse margin, embolium of the hemelytra yellowish with distinct blackish colouration, and abdomen with appressed hairs and large blackish spots, heavy marginal spines on laterotergites III–VI, and elongate light-colored hair.

Etymology: This specific epithet “stephanus” refers to the Institution where this significant research was conducted at the Zoology Research Centre, St. Stephen’s College, Pathanapuram. St. Stephen’s College was founded in 1964 after the name of St. Stephen, one of the first seven ordained Deacons and the first Martyr of Christianity.

Table 1. Comparison among the congeners in the genus *Heleocoris* reported from Kerala, Tamil Nadu, and central states of India.

Taxon	Size	Shape	Head	Pronotum	Scutellum	Hemelytron	Abdomen
<i>H. bergerothi</i>	Length 9.7; width 6.8 mm	Oval	Palely yellow with indistinct brown spots	Colouration same as head, the lateral margins arcuate, the lateral posterior angles acute, the surface densely and finely punctate	Black with yellow apex, very finely granulose, with a yellow v-shaped discal spot	Blackish-brown with largely yellowish embolium	Connexivum, abdomen, and legs entirely pale yellowish
<i>H. vicinus</i>	Length 9–10	Oval	Twice as broad at base between eyes, rounded in front, ochraceous with black speckles	Ochraceous with black speckles, the lateral margins pale ochraceous, distinctly subamplyate, the posterior margin a little paler than the disk	Black with ochraceous apex	Black, lateral margin of corium pale ochraceous with two blackish macular suffusions	Connexivum pale ochraceous with the posterior margins black; abdomen and legs pale ochraceous, left sixth ventral laterotergite is completely flat
<i>H. breviceps</i>	Length 9.5; width 6.5 mm	Oval	Ochraceous; intraocular area marked on each side by two small foveations	Ochraceous	Blackish-brown with the disk narrowly yellow	Hemelytra brownish; embolium ochraceous	Abdomen beneath largely blackish with yellowish apex; legs entirely yellow
<i>H. indicus</i>	Length 8; width 5 mm	Oval	Brown punctures more or less dense forming small ill-defined shadings	Colouration same as head	Dark brown with a median longitudinal line.	Embolium yellowish with a medial brown spot on its inner margin and another spot at its apex	Connexivum yellow, very narrowly darkened on the outer edge of the posterior angles of each segment; underside of body entirely yellowish
<i>H. majusculus</i>	Length 13; width 8.8 mm	Oval	Surface superficially punctate, two impressions, one before the other on each side near the eyes	Very transverse, yellowish, finely granulose, punctured on all the surface, with the anterior margin obtusely sinuate on each side behind the eyes	Entirely black; claval commissure nearly as long as the scutellum	Embolium very much dilated behind	Connexivum yellow, with the posterior angles of the segments a little acuminate; underside of body pitchy brown
<i>H. rotundatus</i>	Length 12; width 9.4 mm	Oval	Thick, finely granulose, brownish-ochraceous, centrally blackish	Brownish-ochraceous much suffused with blackish except on the lateral margin areas which are distinctly rounded	Black with apex ochraceous, strongly and mostly transversely wrinkled; clavalsuture longer than scutellum and apically spotted with ochraceous	Black, thick, finely punctate, lateral margins of embolium ochraceous.	Connexivum ochraceous; under side of the body blackish; legs ochraceous; abdomen beneath with a central longitudinal series of black spots, the lateral margins ochraceous.
<i>H. acutus</i>		Oval		Shape is not perceptibly "flange" with no marginal furrow, the posterior angles acute			
<i>H. elongatus</i>	Length 8; width 4.5 mm	Oval	Clear yellow tint densely punctured with black	Colouration same as head; lateral margins very slightly arcuated nearly straight, the lateral posterior angles rounded, not prominent behind, the posterior edge straight	Brownish; claval suture a little longer than half the length of the scutellum	Embolium little enlarged and largely yellowish and along the outer margin, the yellow margin bisinuated at its inner edge by two brown spots	Connexivum yellow, with a straight transverse brown fascia on the posterior margins of the segments; underside of the body entirely yellow

Taxon	Size	Shape	Head	Pronotum	Scutellum	Hemelytron	Abdomen
<i>H. bengalensis</i>	Length 8 mm	Oval	Posterior margin with brown punctures coalescent and dark punctuation and associated maculation denser	Dark punctuation and associated maculation denser; posterior margin with longitudinal marks		Large yellow patch on embolium.	Abdomen laterally slightly surpassing hemelytra in males; ventral laterotergites II–VII each light anteriorly, dark posteriorly
<i>H. stephanus</i> sp. nov.	Length 9.78; width 6.20 mm	Oval	Densely punctured with black. Single sclerite at posterior corner of eye, directed ventrad, anterior end simple. Goblet shaped black colouration in the dorsal surface	Finely granulose, broad and maximum width at rounded posterolateral corners; tuft of setae in anterior end, transverse sulcus band ending at anterior end of clavus with ventral tuft of setae	Finely granulose, tumescent and elevated above level of wings, sinuated lateral margins.	Blackish-brown, largely yellowish embolium with distinct black colouration	Abdomen ventrally covered with appressed hairs; heavy marginal spines on laterotergites III–VI and brush of elongate light-coloured hairs; legs brownish with setae and bristles.

DISCUSSION

This species was found in the upstream of Kallada River, Kerala, India. The type locality is located in the Western Ghats of the southwestern region of the country. The sampling area of the present investigation was characterized by partially decomposed plant parts. Studies on saucer bugs are scanty in our country, and the sampling area is an undisturbed and unexplored area of Kallada River. There were no detailed recent taxonomical reports on creeping water bugs from the freshwater ecosystems of Kerala. As part of our macroinvertebrate sampling, we explored several ignored parts of the upstream forested area of Kallada River with a rich diversity of creeping water bugs. No additional *Heleocoris* species were collected. Males of this species are most similar to *H. bergrothi* and *H. indicus* based on some external features like coloration pattern, though, these species are clearly distinct from *H. stephanus* in several aspects. Previously, 11 species were reported in India. The present discovery brings the total number of *Heleocoris* species in India to 12.

CONCLUSION

Heleocoris stephanus sp. nov. shows close morphological similarity with *H. bergrothi*, and *H. indicus*; The latter species are differentiated by distinctive morphological and genital features.

REFERENCES

Chandra, K., K.C. Gopi, D.V. Rao, K. Valarmathi & J.R.B. Alfred (2017). Chapter 23: Insecta: Hemiptera, pp. 357–378. In: Basu, S. & K.A. Subramanian (eds.). *Current Status of Freshwater Faunal Diversity in India*. Zoological Survey of India, Kolkata.

Montandon, A.L. (1897). Hemiptera cryptocerata. Fam. Naucoridae.—Sous-fam. Laccocorinae. Verhandlungen der kaiserlich-königlichen zoologisch-botanischen Gesellschaft in Wien 47: 435–454.

Montandon, A. L., (1908). *Heleocoris rotundatus*, La Rivers, Dcc. Pap Bioi. Soc Nevada, 38: 12.

Montandon, A.L. (1910). Espèces nouvelles ou peu connues d'Hydrocorisés de l'Inde. *Bulletin de la Société des Sciences de Bucharest-Roumanie* 19: 652–658.

Polhemus, D.A. & J.T. Polhemus (2013). Guide to the aquatic Heteroptera of Singapore and Peninsular Malaysia. Xi. Infraorder Nepomorpha—families Naucoridae and Aphelocheiridae. *Raffles Bulletin of Zoology* 61(2): 665–686.

Sites, R.W. & A. Vitheerapadit (2011). *Heleocoris* (Heteroptera: Naucoridae: Laccocorinae) of Thailand, with description of a new species. *Zootaxa* 2736(1): 1–16. <https://doi.org/10.1111/zootaxa.2736.1.1>

Stål, C. (1876). Enumeratio Hemipterorum, Bidrag till en förteckning öfver alla hitills kända Hemiptera, jemtens systematicameddelanden. 5. *Kongliga Svenska Vetenskaps-Akademiens Handlingar* 14(4): 1–162.

Xie, T.Y. & G.Q. Liu (2015). The creeping water bugs (Hemiptera: Heteroptera: Naucoridae) of China, with description of a new species. *Zootaxa* 3911(4): 571–580. <https://doi.org/10.1111/zootaxa.3911.4.6>

Mr. Jatishwor Singh Irungbam, Biology Centre CAS, Branišovská, Czech Republic.
Dr. Ian J. Kitching, Natural History Museum, Cromwell Road, UK
Dr. George Mathew, Kerala Forest Research Institute, Peechi, India
Dr. John Noyes, Natural History Museum, London, UK
Dr. Albert G. Orr, Griffith University, Nathan, Australia
Dr. Sameer Padhye, Katholieke Universiteit Leuven, Belgium
Dr. Nancy van der Poorten, Toronto, Canada
Dr. Karen Schnabel, NIWA, Wellington, New Zealand
Dr. R.M. Sharma, (Retd.) Scientist, Zoological Survey of India, Pune, India
Dr. Manju Siliwal, WILD, Coimbatore, Tamil Nadu, India
Dr. G.P. Sinha, Botanical Survey of India, Allahabad, India
Dr. K.A. Subramanian, Zoological Survey of India, New Alipore, Kolkata, India
Dr. P.M. Sureshan, Zoological Survey of India, Kozhikode, Kerala, India
Dr. R. Varatharajan, Manipur University, Imphal, Manipur, India
Dr. Eduard Vives, Museu de Ciències Naturals de Barcelona, Terrassa, Spain
Dr. James Young, Hong Kong Lepidopterists' Society, Hong Kong
Dr. R. Sundararaj, Institute of Wood Science & Technology, Bengaluru, India
Dr. M. Nithyanandan, Environmental Department, La Al Kuwait Real Estate. Co. K.S.C., Kuwait
Dr. Himender Bharti, Punjabi University, Punjab, India
Mr. Purnendu Roy, London, UK
Dr. Saito Motoki, The Butterfly Society of Japan, Tokyo, Japan
Dr. Sanjay Sondhi, TITLI TRUST, Kalpavriksh, Dehradun, India
Dr. Nguyen Thi Phuong Lien, Vietnam Academy of Science and Technology, Hanoi, Vietnam
Dr. Nitin Kulkarni, Tropical Research Institute, Jabalpur, India
Dr. Robin Wen Jiang Ngiam, National Parks Board, Singapore
Dr. Lional Monod, Natural History Museum of Geneva, Genève, Switzerland.
Dr. Asheesh Shivam, Nehru Gram Bharti University, Allahabad, India
Dr. Rosana Moreira da Rocha, Universidade Federal do Paraná, Curitiba, Brasil
Dr. Kurt R. Arnold, North Dakota State University, Saxony, Germany
Dr. James M. Carpenter, American Museum of Natural History, New York, USA
Dr. David M. Claborn, Missouri State University, Springfield, USA
Dr. Karen Schnabel, Marine Biologist, Wellington, New Zealand
Dr. Amazonas Chagas Júnior, Universidade Federal de Mato Grosso, Cuiabá, Brasil
Mr. Monsoon Jyoti Gogoi, Assam University, Silchar, Assam, India
Dr. Heo Chong Chin, Universiti Teknologi MARA (UiTM), Selangor, Malaysia
Dr. R.J. Shiel, University of Adelaide, SA 5005, Australia
Dr. Siddharth Kulkarni, The George Washington University, Washington, USA
Dr. Priyadarshan Dharma Rajan, ATREE, Bengaluru, India
Dr. Phil Alderslade, CSIRO Marine And Atmospheric Research, Hobart, Australia
Dr. John E.N. Veron, Coral Reef Research, Townsville, Australia
Dr. Daniel Whitmore, State Museum of Natural History Stuttgart, Rosenstein, Germany.
Dr. Yu-Feng Hsu, National Taiwan Normal University, Taipei City, Taiwan
Dr. Keith V. Wolfe, Antioch, California, USA
Dr. Siddharth Kulkarni, The Hormiga Lab, The George Washington University, Washington, D.C., USA
Dr. Tomas Ditrich, Faculty of Education, University of South Bohemia in Ceske Budejovice, Czech Republic
Dr. Mihaly Foldvari, Natural History Museum, University of Oslo, Norway
Dr. V.P. Uniyal, Wildlife Institute of India, Dehradun, Uttarakhand 248001, India
Dr. John T.D. Caleb, Zoological Survey of India, Kolkata, West Bengal, India
Dr. Priyadarshan Dharma Rajan, Ashoka Trust for Research in Ecology and the Environment (ATREE), Royal Enclave, Bangalore, Karnataka, India

Fishes

Dr. Neelesh Dahanukar, IISER, Pune, Maharashtra, India
Dr. Topiltzin Contreras MacBeath, Universidad Autónoma del estado de Morelos, México
Dr. Heok Hee Ng, National University of Singapore, Science Drive, Singapore
Dr. Rajeev Raghavan, St. Albert's College, Kochi, Kerala, India
Dr. Robert D. Sluka, Chiltern Gateway Project, A Rocha UK, Southall, Middlesex, UK
Dr. E. Vivekanandan, Central Marine Fisheries Research Institute, Chennai, India
Dr. Davor Zanella, University of Zagreb, Zagreb, Croatia
Dr. A. Biju Kumar, University of Kerala, Thiruvananthapuram, Kerala, India
Dr. Akhilesh K.V., ICAR-Central Marine Fisheries Research Institute, Mumbai Research Centre, Mumbai, Maharashtra, India
Dr. J.A. Johnson, Wildlife Institute of India, Dehradun, Uttarakhand, India
Dr. R. Ravinesh, Gujarat Institute of Desert Ecology, Gujarat, India

Amphibians

Dr. Sushil K. Dutta, Indian Institute of Science, Bengaluru, Karnataka, India
Dr. Annemarie Ohler, Muséum national d'Histoire naturelle, Paris, France

Reptiles

Dr. Gernot Vogel, Heidelberg, Germany
Dr. Raju Vyas, Vadodara, Gujarat, India
Dr. Pritpal S. Soorae, Environment Agency, Abu Dhabi, UAE.
Prof. Dr. Wayne J. Fuller, Near East University, Mersin, Turkey
Prof. Chandrashekher U. Rironker, Goa University, Taleigao Plateau, Goa, India
Dr. S.R. Ganesh, Chennai Snake Park, Chennai, Tamil Nadu, India
Dr. Himansu Sekhar Das, Terrestrial & Marine Biodiversity, Abu Dhabi, UAE

Journal of Threatened Taxa is indexed/abstracted in Bibliography of Systematic Mycology, Biological Abstracts, BIOSIS Previews, CAB Abstracts, EBSCO, Google Scholar, Index Copernicus, Index Fungorum, JournalSeek, National Academy of Agricultural Sciences, NewJour, OCLC WorldCat, SCOPUS, Stanford University Libraries, Virtual Library of Biology, Zoological Records.

NAAS rating (India) 5.64

Birds

Dr. Hem Sagar Baral, Charles Sturt University, NSW Australia
Mr. H. Biju, Coimbatore, Tamil Nadu, India
Dr. Chris Bowden, Royal Society for the Protection of Birds, Sandy, UK
Dr. Priya Davidar, Pondicherry University, Kalapet, Puducherry, India
Dr. J.W. Duckworth, IUCN SSC, Bath, UK
Dr. Rajah Jayopal, SACON, Coimbatore, Tamil Nadu, India
Dr. Rajiv S. Kalsi, M.L.N. College, Yamuna Nagar, Haryana, India
Dr. V. Santharam, Rishi Valley Education Centre, Chittoor Dt., Andhra Pradesh, India
Dr. S. Balachandran, Bombay Natural History Society, Mumbai, India
Mr. J. Praveen, Bengaluru, India
Dr. C. Srinivasulu, Osmania University, Hyderabad, India
Dr. K.S. Gopi Sundar, International Crane Foundation, Baraboo, USA
Dr. Gombobaatar Sundev, Professor of Ornithology, Ulaanbaatar, Mongolia
Prof. Reuven Yosef, International Birding & Research Centre, Eilat, Israel
Dr. Taej Mundkur, Wetlands International, Wageningen, The Netherlands
Dr. Carol Inskip, Bishop Auckland Co., Durham, UK
Dr. Tim Inskip, Bishop Auckland Co., Durham, UK
Dr. V. Gokula, National College, Tiruchirappalli, Tamil Nadu, India
Dr. Arkady Lelej, Russian Academy of Sciences, Vladivostok, Russia
Dr. Simon Dowell, Science Director, Chester Zoo, UK
Dr. Mário Gabriel Santiago dos Santos, Universidade de Trás-os-Montes e Alto Douro, Quinta de Prados, Vila Real, Portugal
Dr. Grant Connette, Smithsonian Institution, Royal, VA, USA
Dr. P.A. Azeez, Coimbatore, Tamil Nadu, India

Mammals

Dr. Giovanni Amori, CNR - Institute of Ecosystem Studies, Rome, Italy
Dr. Anwaruddin Chowdhury, Guwahati, India
Dr. David Mallon, Zoological Society of London, UK
Dr. Shomita Mukherjee, SACON, Coimbatore, Tamil Nadu, India
Dr. Angie Appel, Wild Cat Network, Germany
Dr. P.O. Nameer, Kerala Agricultural University, Thrissur, Kerala, India
Dr. Ian Redmond, UNEP Convention on Migratory Species, Lansdown, UK
Dr. Heidi S. Riddle, Riddle's Elephant and Wildlife Sanctuary, Arkansas, USA
Dr. Karin Schwartz, George Mason University, Fairfax, Virginia.
Dr. Lala A.K. Singh, Bhubaneswar, Orissa, India
Dr. Mewa Singh, Mysore University, Mysore, India
Dr. Paul Racey, University of Exeter, Devon, UK
Dr. Honnavalli N. Kumara, SACON, Anaikatty P.O., Coimbatore, Tamil Nadu, India
Dr. Nishith Dharaiya, HNG University, Patan, Gujarat, India
Dr. Spartaco Gippoliti, Socio Onorario Società Italiana per la Storia della Fauna "Giuseppe Altobello", Rome, Italy
Dr. Justus Joshua, Green Future Foundation, Tiruchirappalli, Tamil Nadu, India
Dr. H. Raghuram, The American College, Madurai, Tamil Nadu, India
Dr. Paul Bates, Harison Institute, Kent, UK
Dr. Jim Sanderson, Small Wild Cat Conservation Foundation, Hartford, USA
Dr. Dan Challender, University of Kent, Canterbury, UK
Dr. David Mallon, Manchester Metropolitan University, Derbyshire, UK
Dr. Brian L. Cypher, California State University-Stanislaus, Bakersfield, CA
Dr. S.S. Talmale, Zoological Survey of India, Pune, Maharashtra, India
Prof. Karan Bahadur Shah, Budhanilkantha Municipality, Kathmandu, Nepal
Dr. Susan Cheyne, Borneo Nature Foundation International, Palangkaraya, Indonesia
Dr. Hemanta Kafley, Wildlife Sciences, Tarleton State University, Texas, USA

Other Disciplines

Dr. Aniruddha Belsare, Columbia MO 65203, USA (Veterinary)
Dr. Mandar S. Paingankar, University of Pune, Pune, Maharashtra, India (Molecular)
Dr. Jack Tordoff, Critical Ecosystem Partnership Fund, Arlington, USA (Communities)
Dr. Ulrike Streicher, University of Oregon, Eugene, USA (Veterinary)
Dr. Hari Balasubramanian, EcoAdvisors, Nova Scotia, Canada (Communities)
Dr. Rayanna Helleni Santos Bezerra, Universidade Federal de Sergipe, São Cristóvão, Brazil
Dr. Jamie R. Wood, Landcare Research, Canterbury, New Zealand
Dr. Wendy Collinson-Jonker, Endangered Wildlife Trust, Gauteng, South Africa
Dr. Rajeshkumar G. Jani, Anand Agricultural University, Anand, Gujarat, India
Dr. O.N. Tiwari, Senior Scientist, ICAR-Indian Agricultural Research Institute (IARI), New Delhi, India
Dr. L.D. Singla, Guru Angad Dev Veterinary and Animal Sciences University, Ludhiana, India
Dr. Rupika S. Rajakaruna, University of Peradeniya, Peradeniya, Sri Lanka
Dr. Bharat Baviskar, Wild-CER, Nagpur, Maharashtra 440013, India

Reviewers 2021–2023

Due to paucity of space, the list of reviewers for 2021–2023 is available online.

The opinions expressed by the authors do not reflect the views of the Journal of Threatened Taxa, Wildlife Information Liaison Development Society, Zoo Outreach Organization, or any of the partners. The journal, the publisher, the host, and the partners are not responsible for the accuracy of the political boundaries shown in the maps by the authors.

Print copies of the Journal are available at cost. Write to:
The Managing Editor, JoTT,
c/o Wildlife Information Liaison Development Society,
43/2 Varadarajulu Nagar, 5th Street West, Ganapathy, Coimbatore,
Tamil Nadu 641006, India
ravi@threatenedtaxa.org

Articles

Spatial assemblage of shorebirds (Aves: Charadriiformes) in an altered wetland of the southern coast of Sri Lanka

– V.N. Mendis, E.J.A.P. Buddhi Priyankara, E.G.D.P. Jayasekara & W.A.D. Mahaulpatha, Pp. 25495–25506

Current conservation status of Bengal Florican *Houbaropsis bengalensis* in Manas National Park, Assam, India

– Miranda Thakur, Jonmani Kalita, Namita Brahma, Koushik Rajbongshi, Kangkanjyoti Bhattacharyya, Amal Chandra Sarmah, Alolika Sinha, Deba Kumar Dutta, Dhritiman Das & Bibhuti Prasad Lahkar, Pp. 25507–25515

Assemblages of frugivorous butterflies in two urban parks in Quezon City, Philippines

– Michael Gabriel A. Itliong, Nikki Heherson A. Dagamac & Jade Aster T. Badon, Pp. 25516–25527

Assessment of the status of *Spodoptera* species (Lepidoptera: Noctuidae: Armyworm) in India through DNA barcoding technique

– Dinesh Nalage, P.S. Kudnar, Tejswini Sontakke, Ishwar Chittapure, Yashdeep Gowda, Shantanu Kharbal & Yashashri Alamwar, Pp. 25528–25535

Taxonomy and distribution of some orthopteran species (Orthoptera: Gryllidae, Trigonidiidae, Acrididae) from northwestern Morocco

– Hanae El Harche, Samiha Kaioua & Dalale Mansouri, Pp. 25536–25544

Impact of root harvest on *Decalepis hamiltonii* Wight & Arn. population across habitats in Savandurga Reserve Forest, Karnataka, India

– M. Sathy Sangeetha, Kaliamoorthy Ravikumar & H.C. Chetan, Pp. 25545–25570

Communications

Rare encounters: Jungle Cat *Felis chaus* Schreber, 1777 (Mammalia: Carnivora: Felidae) in the lower reaches of the Jordan River, Jordan

– Ehab Eid & Mohammad Farid Alayyan, Pp. 25571–25576

Diversity of bird species in Ebpanan Marsh, Maguindanao del Norte, Bangsamoro Autonomous Region in Muslim Mindanao (BARMM), Philippines

– Gindol Rey A. Limbaro, Benito Anthony A. Pingoy & Peter Jan D. de Vera, Pp. 25577–25583

Heleocoris stephanus (Heteroptera: Naucoridae: Laccocorinae), a new species of creeping water bug from Kallada River, Kerala, India

– Dani Benchamin, R. Sreejai & M.S. Arya, Pp. 25584–25589

Incidence and risk factors associated with parasitic infections in captive wild mammals and birds in Indian zoos

– Nikita Das, P.D. Pawar, P.P. Mhase, V.G. Nimbalkar, R.V. Jadhav, V.S. Dhaygude, Gavin Furtado & L.D. Singla, Pp. 25590–25597

Bryophyte diversity of Berinag (Pithoragarh District) in Kumaun Himalaya, Uttarakhand, India

– D. Dhami & P. Chaturvedi, Pp. 25598–25603

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

July 2024 | Vol. 16 | No. 7 | Pages: 25495–25638

Date of Publication: 26 July 2024 (Online & Print)

DOI: 10.11609/jott.2024.16.7.25495-25638

Short Communications

The opportunistic feeding behaviour of *Schistura notostigma* (Teleostei: Nemacheilidae) in tropical mountain streams in Sri Lanka

– J. Bandara, M.P. Gunawardena & R.T.P Jayasuriya, Pp. 25604–25608

First record of *Pieris napi* L. (Lepidoptera: Pieridae) from Kashmir Valley, India

– Firdousa Rasool & Altaf Hussain Mir, Pp. 25609–25612

Reassessment of *Strobilanthes recurva* (Acanthaceae), an endangered plant from Manipur, India

– Rajkumari Jashmi Devi & Biseshwori Thongam, Pp. 25613–25616

New distribution record of Slender Wild Basil *Clinopodium gracile* (Benth.) Kuntze (Lamiaceae: Nepetoideae: Mentheae) for the flora of Himachal Pradesh, India

– Rimjhim Chandra & Mamita Kalita, Pp. 25617–25622

Notes

Rusty-spotted Cat *Prionailurus rubiginosus* (I. Geoffroy Saint-Hilaire, 1831) (Mammalia: Carnivora: Felidae) in the semi-natural subterranean habitat in Karnataka, India

– Shirish Manchi, Goldin Quadros, Dipika Bajpai, Shomita Mukherjee, Suma Haleholi, Mahesh Marennavar, Sangmesh Neeralagi, Prakash Ganiger, Suresh Lamani & Nikhil Kulkarni, Pp. 25623–25626

First record of Scaly-breasted Munia *Lonchura punctulata* (Linnaeus, 1758) (Aves: Passeriformes: Estrildidae) from Kashmir, India

– Shazia Shafayat, Fayaz Ahmad Ahanger, Tariq Ahmad, Bilal A. Bhat & Zakir Hussain Najar, Pp. 25627–25629

First record of *Prosyznskia diatreta* (Simon, 1902) (Araneae: Salticidae) from Gujarat, India

– Manisha P. Patel & Dhruv A. Prajapati, Pp. 25630–25631

Medicago monantha (Fabaceae) and *Euphorbia jodhpurensis* (Euphorbiaceae) as new additions to the flora of Maharashtra State, India

– Praveen V. Kale & Rajendra D. Shinde, Pp. 25632–25636

Book Review

All eyes on the island: A book review of The Great Nicobar Betrayal

– Lakshmi Ravinder Nair, Pp. 25637–25638

Publisher & Host



Threatened Taxa